

成人免疫注射禁忌症的篩選核對表

患者姓名 _____

出生日期 ____/____/____
月 日 年

致患者：下列問題有助於我們確定今天應該為您注射何種疫苗。如果您對所有問題的答案都為「是」，並不一定代表您不應該接種疫苗，而是意指您必須回答額外的問題。如果您對某個問題不太清楚，請要求您的醫護人員說明。

| | 是 | 否 | 不知道 |
|---|--------------------------|--------------------------|--------------------------|
| 1. 您今天是否感到不舒服？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. 您是否對藥物食物、任何疫苗成份、或乳膠過敏？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. 您是否曾經在接種疫苗後出現過嚴重反應？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. 您是否患有心臟病、肺病，氣喘、腎臟病、代謝性疾病 (如糖尿病)、貧血、或其他血液障礙的長期疾病？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 5. 您是否患有癌症、白血病、HIV/愛滋病或有任何其他免疫系統問題？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 6. 在過去三個月內您是否曾服用減弱免疫力的藥物如強的松、其他類固醇，或是抗癌藥物；治療風濕性關節炎、克隆氏症或牛皮癬的藥物；或是接受放射線治療？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 7. 您是否患有癲癇或大腦或神經問題？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 8. 在過去一年，您是否接受過輸血或使用血液製品，或使用過稱為免疫 (gamma) 球蛋白的藥物或抗病毒藥物？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 9. 對於女性患者：您是否懷孕，或在下個月有懷孕的可能？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 10. 您在過去4週中是否注射過疫苗？ | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

填表人 _____ 日期 _____

表格審核人 _____ 日期 _____

您是否攜帶您的免疫注射紀錄卡？ 是 否

擁有一份個人疫苗記錄非常重要。如果您沒有該記錄，請向您的醫療保健提供者索取。請妥善保管記錄，每次就診時隨身攜帶。確定您的醫療保健者將您的所有免疫注射記錄在上面。

Information for Healthcare Professionals about the Screening Checklist for Contraindications to Vaccines for Adults

Are you interested in knowing why we included a certain question on the screening checklist? If so, read the information below. If you want to find out even more, consult the references listed at the end.

1. Are you sick today? [all vaccines]

There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events.¹ However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. Mild illnesses (such as upper respiratory infections or diarrhea) are NOT contraindications to vaccination. Do not withhold vaccination if a person is taking antibiotics.

2. Do you have allergies to medications, food, a vaccine component, or latex? [all vaccines]

An anaphylactic reaction to latex is a contraindication to vaccines that contain latex as a component or as part of the packaging (e.g., vial stoppers, prefilled syringe plungers, prefilled syringe caps). If a person has anaphylaxis after eating gelatin, do not administer vaccines containing gelatin. A local reaction to a prior vaccine dose or vaccine component, including latex, is not a contraindication to a subsequent dose or vaccine containing that component. For information on vaccines supplied in vials or syringes containing latex, see reference 2; for an extensive list of vaccine components, see reference 3.

People with egg allergy of any severity can receive any IIV, RIV, or LAIV that is otherwise appropriate for the patient's age and health status. The safety of LAIV in egg allergic people has not been established. For people with a history of severe allergic reaction to egg involving any symptom other than hives (e.g., angioedema, respiratory distress), or who required epinephrine or another emergency medical intervention, the vaccine should be administered in a medical setting, such as a clinic, health department, or physician office. Vaccine administration should be supervised by a healthcare provider who is able to recognize and manage severe allergic conditions.⁴

3. Have you ever had a serious reaction after receiving a vaccination? [all vaccines]

History of anaphylactic reaction (see question 2) to a previous dose of vaccine or vaccine component is a contraindication for subsequent doses.¹ Under normal circumstances, vaccines are deferred when a precaution is present. However, situations may arise when the benefit outweighs the risk (e.g., during a community pertussis outbreak).

4. Do you have a long-term health problem with heart disease, lung disease, asthma, kidney disease, metabolic disease (e.g., diabetes), anemia, or other blood disorder? [MMR, LAIV]

A history of thrombocytopenia or thrombocytopenic purpura is a precaution to MMR vaccine. These conditions, including asthma in adults, should be considered precautions for the use of LAIV.

5. Do you have cancer, leukemia, HIV/AIDS, or any other immune system problem? [LAIV, MMR, VAR, ZVL]

Live virus vaccines (e.g., LAIV, measles-mumps-rubella [MMR], varicella [VAR], zoster vaccine live [ZVL]) are usually contraindicated in immunocompromised people. However, there are exceptions. For example, MMR vaccine is recommended and varicella vaccine should be considered for adults with CD4+ T-lymphocyte counts of greater than or equal to 200 cells/ μ L. Immunosuppressed people should not receive LAIV. For details, consult the ACIP recommendations.^{4,5,6}

6. In the past 3 months, have you taken medications that affect your immune system, such as cortisone, prednisone, other steroids, or anticancer drugs; drugs for the treatment of rheumatoid arthritis, Crohn's disease, or psoriasis; or have you had radiation treatments? [LAIV, MMR, VAR, ZVL]

Live virus vaccines (e.g., LAIV, MMR, VAR, ZVL) should be postponed until after chemotherapy or long-term high-dose steroid therapy has ended. For details and length of time to postpone, consult the ACIP statement.^{1,5} Some immune mediator and immune modulator drugs (especially the anti-tumor necrosis factor agents adalimumab, infliximab, etanercept, golimumab, and certolizumab pegol) may be immunosuppressive. The use of live vaccines should be avoided

in persons taking these drugs (see www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html). To find specific vaccination schedules for stem cell transplant (bone marrow transplant) patients, see reference 7. LAIV can be given only to healthy non-pregnant people ages 2 through 49 years.

7. Have you had a seizure or a brain or other nervous system problem? [influenza, Td/Tdap]

Tdap is contraindicated in people who have a history of encephalopathy within 7 days following DTP/DTaP. An unstable progressive neurologic problem is a precaution to the use of Tdap. For people with stable neurologic disorders (including seizures) unrelated to vaccination, or for people with a family history of seizure, vaccinate as usual. A history of Guillain-Barré syndrome (GBS) is a consideration with the following: 1) Td/Tdap: if GBS has occurred within 6 weeks of a tetanus-toxoid vaccine and decision is made to continue vaccination, give Tdap instead of Td if no history of prior Tdap; 2) Influenza vaccine (IIV/LAIV): if GBS has occurred within 6 weeks of a prior influenza vaccine, vaccinate with IIV if at increased risk for severe influenza complications.

8. During the past year, have you received a transfusion of blood or blood products, or been given immune (gamma) globulin or an antiviral drug? [MMR, VAR]

Certain live virus vaccines (e.g., MMR, VAR) may need to be deferred, depending on several variables. Consult the most current ACIP recommendations for current information on intervals between antiviral drugs, immune globulin or blood product administration and live virus vaccines.¹

9. For women: Are you pregnant or is there a chance you could become pregnant during the next month? [HPV, IPV, MMR, LAIV, VAR, ZVL]

Live virus vaccines (e.g., MMR, VAR, ZVL, LAIV) are contraindicated one month before and during pregnancy because of the theoretical risk of virus transmission to the fetus. Sexually active women in their childbearing years who receive live virus vaccines should be instructed to avoid pregnancy for one month following receipt of the vaccine. On theoretical grounds, inactivated poliovirus vaccine should not be given during pregnancy; however, it may be given if risk of exposure is imminent and immediate protection is needed (e.g., travel to endemic areas). Inactivated influenza vaccine and Tdap are both recommended during pregnancy. Both vaccines may be given at any time during pregnancy but the preferred time for Tdap administration is at 27–36 weeks' gestation. HPV vaccine is not recommended during pregnancy.^{1,4,5,6,8,9}

10. Have you received any vaccinations in the past 4 weeks? [LAIV, MMR, VAR, yellow fever, ZVL]

People who were given either LAIV or an injectable live virus vaccine (e.g., MMR, VAR, ZVL, yellow fever) should wait 28 days before receiving another vaccination of this type (30 days for yellow fever). Inactivated vaccines may be given at any spacing interval if they are not administered simultaneously.

REFERENCES

1. CDC. General best practice guidelines for immunization. Best Practices Guidance of the Advisory Committee on Immunization Practices (ACIP) at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html.
2. Latex in Vaccine Packaging: www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/latex-table.pdf.
3. Table of Vaccine Components: www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf.
4. CDC. Prevention and control of seasonal influenza with vaccines: Recommendations of the Advisory Committee on Immunization Practices – United States, . . . Access links to current ACIP recommendations at www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/flu.html
5. CDC. Measles, mumps, and rubella – vaccine use and strategies for elimination of measles, rubella, and congenital rubella syndrome and control of mumps. *MMWR* 1998; 47 (RR-8).
6. CDC. Prevention of varicella: Recommendations of the Advisory Committee on Immunization Practices. *MMWR* 2007; 56 (RR-4).
7. Tomblin M, Einsele H, et al. Guidelines for preventing infectious complications among hematopoietic stem cell transplant recipients: a global perspective. *Biol Blood Marrow Transplant* 15:1143–1238; 2009 at www.cdc.gov/vaccines/pubs/hemato-cell-transplants.htm.
8. CDC. Notice to readers: Revised ACIP recommendation for avoiding pregnancy after receiving a rubella-containing vaccine. *MMWR* 2001; 50 (49).
9. CDC. Updated recommendations for use of tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccine (Tdap) in pregnant women: Recommendations of the ACIP. *MMWR* 2012; 62 (7):131–4.