

# VACCINATE ADULTS!

from the Immunization Action Coalition — [www.immunize.org](http://www.immunize.org)

## What's In This Issue

ACIP Votes to Update Recommendations .....	1
Ask the Experts: CDC Answers Your Questions ...	1
Vaccine Highlights .....	4
New! MenB Vaccine: CDC Answers Questions ...	5
Updated! MenB Vaccine Recommendations by Age and Risk Factor .....	7
Updated! Standing Orders for MenB Vaccine ...	8
Updated! MenACWY Vaccine Recommendations by Age and Risk Factor .....	9
Standing Orders for MenACWY Vaccines .....	10
Resources to Help Improve Your MenACWY Dose #2 Coverage for 16-Year-Olds .....	11
Updated! Meningococcal Q&As – for Patients ..	12
Meningococcal Handout to Give to Patients .....	13
Which Vaccines Do I Need Today? A Screening Form for Adults .....	14
Updated! IAC's Temperature Logs .....	15
Use this Checklist to Protect Your Vaccines .....	16
Vaccine Handling Tips .....	17
Products You Can Purchase from IAC .....	18
Please Donate to IAC! .....	19
IAC's Immunization Resources Order Form .....	20

## Ask the Experts

The Immunization Action Coalition extends thanks to our experts, medical officer Andrew T. Kroger, MD, MPH, and nurse educator Donna L. Weaver, RN, MN, both with the National Center for Immunization and Respiratory Diseases at the Centers for Disease Control and Prevention (CDC).

### HPV vaccine

#### What is the new HPV vaccine schedule recommendation?

In October 2016, the Advisory Committee on Immunization Practices (ACIP) voted to recommend a routine 2-dose HPV vaccine schedule for adolescents who start the vaccination series before the 15th birthday. The two doses should be separated by 6–12 months (the minimum interval between doses is 5 months). A 3-dose schedule continues to be recommended for people who start the series on or after the 15th birthday and for people

## ACIP Votes to Update Recommendations for HPV, Tdap, MenB, and HepB Vaccines

On October 19–20, CDC's Advisory Committee on Immunization Practices (ACIP) met in Atlanta and voted to update several of its existing vaccine recommendations. Some of the changes are described below.

### Human Papillomavirus (HPV) Vaccine

ACIP voted to change the HPV vaccination schedule from a 3-dose to a 2-dose series for adolescents who begin the HPV series at 9 through 14 years of age, regardless of age at series completion. Those who start the series later, at 15 through 26 years of age, or who are immunocompromised, will continue to need 3 doses.

The 9vHPV vaccine (HPV9, Gardasil 9, Merck) will soon be the only HPV vaccine available in the U.S. As of October 2016, Merck is distributing only HPV9, and supplies of 2vHPV (Cervarix, GSK) in the U.S. are now depleted. HPV9 may be used to complete a series begun with 4vHPV (HPV4, Gardasil, Merck) or 2vHPV.

### Meningococcal Serogroup B Vaccine

Bexsero (MenB-4C, GSK) has previously been recommended by ACIP for use as a 2-dose series for high-risk individuals and in outbreak settings, and may also be administered to healthy individuals age 16 through 23 years. In April, FDA approved a label change giving MenB-FHbp (Trumenba, Pfizer) as either a 2-dose (0, 6 months) or 3-dose

(0, 1–2, 6 months) series. ACIP voted to recommend that healthcare providers who use Trumenba continue to use the 3-dose series when vaccinating people at increased risk of meningococcal serogroup B disease (e.g., people with persistent complement component deficiencies or anatomical or functional asplenia) or during serogroup B outbreaks. The 2-dose series of Trumenba can be used for routine vaccination for healthy people age 16 through 23 years.

### Tdap Vaccine

Previous ACIP recommendations called for prenatal care providers to vaccinate all pregnant women with Tdap vaccine during each pregnancy with optimal timing for this dose designated between 27 and 36 weeks gestation. In October, ACIP voted to recommend administering Tdap vaccination early in the 27- through 36-week "window" to maximize passive antibody transfer to the infant. The new recommendations also clarify that children age 7 through 10 years who receive Tdap as part of a catch-up series may be given an additional Tdap for the routinely recommended adolescent dose at 11–12 years of age.

### Hepatitis B Vaccine

ACIP voted to approve a new guidance document that consolidates all previously published recommendations into a comprehensive statement. ♦

with certain immunocompromising conditions (such as cancer, HIV infection, or taking immunosuppressive drugs). An updated ACIP statement was published on December 15, and is available at [www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6549.pdf](http://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6549.pdf), pages 1405–8.

#### Has ACIP expressed a preference for the 2-dose over the 3-dose schedule for adolescents 9 through 14 years of age?

Yes. ACIP recommends the 2-dose schedule for people starting the HPV vaccination series before the 15th birthday, as long as they are immunocompetent.

#### Does the 2-dose HPV vaccine schedule need to be completed with the same vaccine, or can it include different vaccines (such as bivalent or quadrivalent vaccine)?

The 2-dose schedule can be completed with any combination of HPV vaccine brands as long as dose #1 was given before age 15 years. Dose #2 should be administered 6–12 months after dose #1.

#### If dose #1 of HPV vaccine was given before the 15th birthday and it has been more than a year since that dose was given, would the series be complete with just one additional dose?

Yes. Adolescents and adults who started the HPV vaccine series prior to the 15th birthday and who are not immunocompromised are considered to be adequately vaccinated with just one additional dose of HPV vaccine.

#### We have adolescents in our practice who have received the first 2 doses of the HPV

Ask the Experts...continued on page 2 ►

### Immunization questions?

- Email [nipinfo@cdc.gov](mailto:nipinfo@cdc.gov)
- Call your state health department (phone numbers at [www.immunize.org/coordinators](http://www.immunize.org/coordinators))

## Vaccinate Adults

online at [www.immunize.org/va](http://www.immunize.org/va)  
**Immunization Action Coalition**

2550 University Ave. W., Suite 415 North  
Saint Paul, MN 55114  
Phone: (651) 647-9009  
Email: [admin@immunize.org](mailto:admin@immunize.org)  
Websites: [www.immunize.org](http://www.immunize.org)  
[www.vaccineinformation.org](http://www.vaccineinformation.org)  
[www.immunizationcoalitions.org](http://www.immunizationcoalitions.org)  
[www.give2mcv4.org](http://www.give2mcv4.org)

*Vaccinate Adults* is a publication of the Immunization Action Coalition (IAC) for healthcare professionals. Content is reviewed by the Centers for Disease Control and Prevention (CDC) for technical accuracy. This publication is supported in part by CDC Grant No. 6NH23IP922550. Content is solely the responsibility of IAC and does not necessarily represent the official views of CDC. ISSN 1526-1824.

### Publication Staff

Editor: Deborah L. Wexler, MD  
Associate Editors: William L. Atkinson, MD, MPH; Diane C. Peterson  
Consulting Editors: Teresa Anderson, DDS, MPH; Marian Deegan, JD

### IAC Staff

Chief Strategy Officer:  
L.J. (Litjen) Tan, MS, PhD  
Associate Director for Research:  
Sharon G. Humiston, MD, MPH  
Coordinator for Public Health:  
Laurel Wood, MPA  
Nurse Consultant: Pat Vranesich, RN, BSN  
Coordinator for Hepatitis B Projects:  
Lynn Pollock, RN, MSN  
Perinatal Hepatitis B Consultant:  
Beth Rowe-West, BSN  
Policy Consultant: Sarah R. Landry, MA  
Senior Admin. for Grants and Leadership:  
Julie Murphy, MA  
Senior Project Manager: Robin VanOss  
Operations Manager: Casey Pauly  
Project Administrator: Chrystal Mann

IAC publishes a free email news service (*IAC Express*) and two free periodicals (*Vaccinate Adults* and *Needle Tips*). To subscribe, go to [www.immunize.org/subscribe](http://www.immunize.org/subscribe).

IAC, a 501(c)(3) charitable organization, publishes practical immunization information for healthcare professionals to help increase immunization rates and prevent disease.

### The Immunization Action Coalition is also supported by

Pfizer Inc., Merck Sharp & Dohme Corp., GlaxoSmithKline, Sanofi Pasteur, AstraZeneca, Seqirus, Physicians' Alliance of America, Besse Medical, American Pharmacists Association, Mark and Muriel Wexler Foundation, Herbert and Jeanne Mayer Foundation, and many other generous donors.

IAC maintains strict editorial independence in its publications.

### IAC Board of Directors

Stephanie L. Jakim, MD  
*Olmsted Medical Center*  
Sheila M. Specker, MD  
*University of Minnesota*  
Debra A. Strodthoff, MD  
*Amery Regional Medical Center*  
Deborah L. Wexler, MD  
*Immunization Action Coalition*

## Ask the Experts...continued from page 1

**series 1 or 2 months apart according to the 3-dose schedule. Can we consider their HPV vaccine series to be complete or do we need to give these patients a third dose?**

People who have received 2 doses of HPV vaccine separated by less than 5 months should receive a third dose 6–12 months after dose #1 and at least 12 weeks after dose #2.

**Will the 2-dose recommendation be retroactive for children and teens vaccinated prior to 2016?**

Yes. Any person who ever received 2 doses of any combination of HPV vaccines can be considered fully vaccinated if dose #1 was given before the 15th birthday and the 2 doses were separated by at least 5 months.

## MenACWY vaccine

**Please review the new recommendations for use of MenACWY vaccine in people with human immunodeficiency virus (HIV) infection.**

A growing body of evidence supports an increased risk for meningococcal disease in HIV-infected people. The Advisory Committee on Immunization Practices (ACIP) recommends that all HIV-infected people 2 months of age and older should routinely receive an age-appropriate MenACWY vaccine (Menactra, Sanofi Pasteur; Menveo, GSK).

People age 2 years and older with HIV infection who have not been previously vaccinated should receive a 2-dose primary series of MenACWY vaccine (doses separated by 8–12 weeks). People with HIV infection who have previously received one dose of MenACWY should receive a second dose at the earliest opportunity (at least 8 weeks after the previous dose). Adolescents and adults should receive a booster dose of MenACWY vaccine every 5 years throughout life.

**I have an HIV-positive 64-year-old patient who received MenACWY vaccine last week. Was this the correct vaccine for this patient or should he have gotten meningococcal polysaccharide vaccine (MPSV4, Sanofi Pasteur) due to his age? Also, should this patient get another dose in 2 months?**

MenACWY was the correct vaccine in this situation. The 2013 ACIP recommendations on MenACWY vaccination recommend the use of meningococcal conjugate vaccine in adults age 56 years and older who

were vaccinated previously with MenACWY and now need revaccination, or are recommended to receive multiple doses. A person of this age with HIV infection should receive 2 doses of MenACWY separated by 8–12 weeks. Both MenACWY vaccines are licensed for use in people through age 55 years, which means that the use of these vaccines in people age 56 and older is off-label but recommended by ACIP.

## MenB vaccine

**Which individuals in risk groups are recommended to be vaccinated against meningococcal serogroup B disease?**

CDC's Advisory Committee on Immunization Practices (ACIP) recommends routine MenB vaccination of the following individuals in certain risk groups:

- People age 10 years and older who have functional or anatomic asplenia
- People age 10 years and older who have persistent complement component deficiency, including people taking eculizumab (Soliris)
- People age 10 years and older who are at risk during an outbreak caused by a vaccine serogroup, such as on a college campus
- Microbiologists who work with meningococcus bacteria in a laboratory

Both MenB vaccines are licensed for use in people through age 25 years, which means that the use of these vaccines in people age 26 and older is off-label but recommended by ACIP.

**Which individuals are recommended to be vaccinated against meningococcal serogroup B disease who are not in risk groups?**

ACIP recommends that a MenB vaccine series (Bexsero, MenB-4C, GSK; Trumenba, MenB-FHbp, Pfizer) may be administered to people 16 through 23 years of age with a preferred age of vaccination of 16 through 18 years. This Category B recommendation gives clinicians an opportunity to discuss the value of MenB vaccination with their patients and to make a decision together about the individual's need or desire for the vaccine based on risks, benefits, and wish for protection from the disease. Because it is a Category B recommendation, MenB vaccination is covered by the Vaccines for Children Program for anyone who is eligible. Under the Affordable Care Act, private insurance must also cover the costs of both Category A and B recommended vaccines.

Ask the Experts...continued on page 3 ►

Subscribe to *IAC Express*, the Immunization Action Coalition's e-news and information service at [www.immunize.org/subscribe](http://www.immunize.org/subscribe)

DISCLAIMER: *Vaccinate Adults!* is available to all readers free of charge. Some of the information in this issue is supplied to us by the Centers for Disease Control and Prevention in Atlanta, Georgia, and some information is supplied by third-party sources. The Immunization Action Coalition (IAC) has used its best efforts to accurately publish all of this information, but IAC cannot guarantee that the original information as supplied by others is correct or complete, or that it has been accurately published. Some of the information in this issue is created or compiled by IAC. All of the information in this issue is of a time-critical nature, and we cannot guarantee that some of the information is not now outdated, inaccurate, or incomplete. IAC cannot guarantee that reliance on the information in this issue will cause no injury. Before you rely on the information in this issue, you should first independently verify its current accuracy and completeness. IAC is not licensed to practice medicine or pharmacology, and the providing of the information in this issue does not constitute such practice. Any claim against IAC must be submitted to binding arbitration under the auspices of the American Arbitration Association in St. Paul, Minnesota.

IAC's "Ask the Experts" team from the Centers for Disease Control and Prevention



Andrew T. Kroger, MD, MPH



Donna L. Weaver, RN, MN

Ask the Experts...continued from page 2

**What is the new schedule for Trumenba MenB vaccine?**

The Food and Drug Administration approved a 2-dose schedule for Trumenba in April 2016. At its October 2016 meeting, ACIP voted to recommend a 2-dose schedule of Trumenba for people not at increased risk of MenB (for example, healthy adolescents). The two doses should be administered at least 6 months apart. ACIP recommends that people at increased risk of MenB disease (complement component deficiency, functional or anatomic asplenia, at risk during an outbreak of meningococcal B disease, and certain microbiologists) receive a 3-dose Trumenba series with dose #2 and dose #3 administered 2 and 6 months after dose #1.

The schedule for Bexsero has not changed. Bexsero is a 2-dose series with dose #2 given at least 1 month after dose #1.

**Should college students be vaccinated against meningococcal B disease?**

Although several small meningococcal serogroup B disease outbreaks have occurred on college campuses since 2013, college students in general are not at higher risk of meningococcal B disease than people of the same age who are not college students. Consequently, ACIP does not routinely recommend MenB vaccination for college students. However, college students may choose to receive MenB vaccine to reduce their risk of serogroup B meningococcal disease.

**Stay current with FREE subscriptions**

The Immunization Action Coalition's 2 periodicals, *Vaccinate Adults* and *Needle Tips*, and our email news service, *IAC Express*, are packed with up-to-date information.

**Subscribe to all 3 free publications in one place. It's simple! Go to**

[www.immunize.org/subscribe](http://www.immunize.org/subscribe)

**ACIP recommendations for MenB vaccine say the vaccine will provide "short term protection." What does "short term protection" mean?**

MenB vaccines were approved based on the serologic response to the vaccine. No data are available on vaccine effectiveness against clinical disease or duration of protection against clinical disease. Short term protection refers to the known duration of the antibody response. Available data indicate that a protective antibody level should persist in most recipients for 24–48 months after vaccination. This issue will continue to be monitored. For more information, see the ACIP recommendations at [www.cdc.gov/mmwr/pdf/wk/mm6441.pdf](http://www.cdc.gov/mmwr/pdf/wk/mm6441.pdf), pages 1171–5.

**Can the MenB series be completed with a different MenB brand from the one the series was begun with?**

No. You may not switch MenB vaccines in order to complete a series. The series must be started and completed with the same MenB brand.

**Can meningococcal conjugate (MenACWY) and MenB vaccines be given at the same visit?**

Yes. Meningococcal conjugate and MenB vaccines can be given at the same visit or at any time before or after the other.

**Which groups of patients should receive a booster dose of MenB vaccine after completion of the series?**

ACIP does not currently recommend booster doses of MenB vaccine for any group. ♦

To find more than **1,000** Ask the Experts Q&As answered by CDC experts, visit [www.immunize.org/askexperts](http://www.immunize.org/askexperts)

Additional Ask the Experts Q&As are included in IAC's free weekly e-newsletter, *IAC Express*. Subscribe at [www.immunize.org/subscribe](http://www.immunize.org/subscribe).

**Vaccinate Adults correction policy**

If you find an error, please notify us immediately by sending an email message to [admin@immunize.org](mailto:admin@immunize.org). We publish notification of significant errors in our email announcement service, *IAC Express*. Be sure you're signed up for this service. To subscribe, visit [www.immunize.org/subscribe](http://www.immunize.org/subscribe).

**Advisory Board**

**Liaisons from Organizations**

- Bernadette A. Albanese, MD, MPH**  
Council of State & Territorial Epidemiologists
  - Stephen L. Cochi, MD, MPH**  
Nat'l Ctr. for Immun. & Resp. Diseases, CDC
  - Bruce Gellin, MD, MPH**  
National Vaccine Program Office, DHHS
  - Neal A. Halsey, MD**  
Institute for Vaccine Safety, Johns Hopkins Univ.
  - Claire Hannan, MPH**  
Association of Immunization Managers
  - Carol E. Hayes, CNM, MN, MPH**  
American College of Nurse-Midwives
  - Gregory James, DO, MPH, FACOFP**  
American Osteopathic Association
  - Samuel L. Katz, MD**  
Pediatric Infectious Diseases Society
  - Elyse Olshen Kharbanda, MD, MPH**  
Society for Adolescent Health and Medicine
  - Marie-Michele Leger, MPH, PA-C**  
American Academy of Physician Assistants
  - Kimberly Martin**  
Assn. of State & Territorial Health Officials
  - Lisa M. McKeown, MPH**  
Nat'l. Assn. of County & City Health Officials
  - Kathleen M. Neuzil, MD, MPH**  
American College of Physicians
  - Paul A. Offit, MD**  
Vaccine Education Ctr., Children's Hosp. of Phila.
  - Walter A. Orenstein, MD**  
Emory Vaccine Center, Emory University
  - Mitchel C. Rothholz, RPh, MBA**  
American Pharmacists Association
  - Thomas N. Saari, MD**  
American Academy of Pediatrics
  - Margot L. Savoy, MD, MPH**  
American Academy of Family Physicians
  - William Schaffner, MD**  
Infectious Diseases Society of America
  - Anne Schuchat, MD**  
Centers for Disease Control and Prevention
  - Rhoda Sperling, MD**  
Amer. College of Obstetricians & Gynecologists
  - Thomas E. Stenvig, RN, PhD**  
American Nurses Association
  - Ann S. Taub, MA, CPNP**  
National Assn. of Pediatric Nurse Practitioners
  - John W. Ward, MD**  
Division of Viral Hepatitis, NCHHSTP, CDC
  - Patricia N. Whitley-Williams, MD, MPH**  
National Medical Association
  - Walter W. Williams, MD, MPH**  
Nat'l Ctr. for Immun. & Resp. Diseases, CDC
- Individuals**
- Hie-Won L. Hann, MD**  
Jefferson Medical College, Philadelphia
  - Mark A. Kane, MD, MPH**  
Seattle, Washington
  - Edgar K. Marcuse, MD, MPH**  
University of Washington School of Medicine
  - Harold S. Margolis, MD**  
Taos, New Mexico
  - Brian J. McMahon, MD**  
Alaska Native Medical Center, Anchorage
  - Stanley A. Plotkin, MD**  
Vaxconsult.com
  - Gregory A. Poland, MD**  
Mayo Clinic
  - Sarah Jane Schwarzenberg, MD**  
University of Minnesota
  - Coleman I. Smith, MD**  
Minnesota Gastroenterology, Minneapolis
  - Richard K. Zimmerman, MD, MPH**  
University of Pittsburgh

# Vaccine Highlights

## Recommendations, schedules, and more

*Editor's note: The information in Vaccine Highlights is current as of December 15, 2016.*

### Next ACIP meetings

The Advisory Committee on Immunization Practices (ACIP) is comprised of 15 national experts who advise CDC on the appropriate use of vaccines.

ACIP meets three times a year in Atlanta; meetings are open to the public and viewable online via live webcast. The next meetings will be held on Feb. 22–23 and June 21–22, 2017. For more information, visit [www.cdc.gov/vaccines/acip](http://www.cdc.gov/vaccines/acip).

ACIP periodically issues recommendations on the use of vaccines; they are published and readily available in the *Morbidity and Mortality Weekly Report (MMWR)*. Clinicians who vaccinate should have a current set for reference. Here are sources:

- Download from IAC's website: [www.immunize.org/acip](http://www.immunize.org/acip)
- Download from CDC's website: [www.cdc.gov/vaccines/hcp/acip-recs](http://www.cdc.gov/vaccines/hcp/acip-recs)

In addition, extensive information on ACIP meetings is available at [www.cdc.gov/vaccines/acip/meetings/index.html](http://www.cdc.gov/vaccines/acip/meetings/index.html).

### HPV vaccine news

On December 15, CDC published "Use of a 2-Dose Schedule for Human Papillomavirus Vaccination – Updated Recommendations of the Advisory Committee on Immunization Practices." It is available

at [www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6549.pdf](http://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6549.pdf), pages 1405–8.

On October 7, the Food and Drug Administration approved a 2-dose schedule for Gardasil 9 HPV vaccine (Merck) for people 9 through 14 years of age. The approval was based on a clinical trial that demonstrated a non-inferior response to 2 doses of Gardasil 9 among girls and boys 9 through 14 years of age compared to a 3-dose schedule among women 16 through 26 years of age. The revised package insert and Summary Basis for Regulatory Action is available on the FDA website at [www.fda.gov/BiologicsBloodVaccines/Vaccines/Approved-Products/ucm426445.htm](http://www.fda.gov/BiologicsBloodVaccines/Vaccines/Approved-Products/ucm426445.htm)

On December 2, CDC released an updated human papillomavirus (HPV) vaccine VIS. It is similar to the previous 9-valent HPV vaccine VIS, except that it contains information about the recently approved 2-dose schedule. Providers are encouraged to begin using the new VIS now but may use up stocks of the previous version, especially for patients still using the 3-dose schedule.

The new VIS no longer has "Gardasil-9" in its title because the other two HPV vaccines (Cervarix and quadrivalent Gardasil) are no longer distributed in the U.S. The last doses of Cervarix have already expired, and the last doses of Gardasil will expire in May 2017. At that point Gardasil-9 will be the only HPV vaccine available in the U.S. and this will be the only VIS. The new VIS is available on the on the IAC VIS website at [www.immunize.org/vis/vis\\_hpv\\_gardasil.asp](http://www.immunize.org/vis/vis_hpv_gardasil.asp). Translations of the HPV9 VIS are acceptable to use until new translations become available.

## IAC EXPRESS

Get weekly updates on vaccine information while it's still news!

All the news we publish in "Vaccine Highlights" will be sent by email to you every Wednesday. Free!

To sign up for *IAC Express* – and any of our other free publications – visit

[www.immunize.org/subscribe](http://www.immunize.org/subscribe)

### MenACWY vaccine news

On November 4, CDC published "Recommendations for Use of Meningococcal Conjugate Vaccines in HIV-Infected Persons – ACIP, 2016" in *MMWR*. The document is available at [www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6543.pdf](http://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6543.pdf), pages 1189–94. Routine meningococcal conjugate (MenACWY) vaccination is now recommended for all HIV-infected people age 2 months and older. HIV-infected adolescents and adults should receive 2 doses of MenACWY separated by 8 weeks followed by a booster dose every 5 years throughout their life. ♦

### Current VIS dates

Check the dates on your supply of Vaccine Information Statements (VISs). If they are out of date, obtain the most up-to-date versions as well as VIS translations in more than 30 languages at [www.immunize.org/vis](http://www.immunize.org/vis).

Adenovirus .....	6/11/14	MMRV .....	5/21/10
Anthrax .....	3/10/10	Multi-vaccine .....	11/5/15
Chickenpox .....	3/13/08	PCV13 .....	11/5/15
DTaP .....	5/17/07	PPSV .....	4/24/15
Hib .....	4/2/15	Polio .....	7/20/16
Hepatitis A .....	7/20/16	Rabies .....	10/6/09
Hepatitis B .....	7/20/16	Rotavirus .....	4/15/15
HPV .....	12/2/16	Shingles .....	10/6/09
Influenza .....	8/7/15	Td .....	2/24/15
Japanese enceph...	1/24/14	Tdap .....	2/24/15
MCV4/MPSV4 .....	3/31/16	Typhoid .....	5/29/12
MenB .....	8/9/16	Yellow fever .....	3/30/11
MMR .....	4/20/12		

For a ready-to-print version of this table for posting in your practice, go to [www.immunize.org/catg.d/p2029.pdf](http://www.immunize.org/catg.d/p2029.pdf).

## Apply for IAC's Influenza Vaccination Honor Roll

Join more than 500 healthcare settings already honored!



This honor roll recognizes healthcare settings that have implemented mandatory vaccination policies for healthcare personnel (HCP).

To find the healthcare settings listed by state, visit [www.immunize.org/honor-roll/influenza-mandates/honorees.asp](http://www.immunize.org/honor-roll/influenza-mandates/honorees.asp)

To read position statements supporting mandatory HCP vaccination from leading healthcare organizations and professional medical societies or to apply, visit [www.immunize.org/honor-roll/influenza-mandates](http://www.immunize.org/honor-roll/influenza-mandates)

# Meningococcal B Vaccine: Q&A

## CDC Answers Your Questions

*Experts from the National Center for Immunization and Respiratory Diseases at the Centers for Disease Control and Prevention answer your questions about meningococcal serogroup B (MenB) vaccine.*

### **Which meningococcal vaccines are available in the United States?**

Since 2005, two types of meningococcal vaccines have been available in the United States that protect against meningococcal serogroups A, C, W, and Y: 1) meningococcal polysaccharide vaccine (MPSV4, Menomune, Sanofi Pasteur) which is made up of polysaccharide (sugar molecules) from the surface of the meningococcal bacteria; and 2) meningococcal conjugate vaccines (MenACWY, Menactra, Sanofi Pasteur; Menveo, GSK) in which the polysaccharide is chemically bonded (“conjugated”) to a protein to produce better protection.

More recently, two vaccines have become available that offer protection from meningococcal serogroup B disease (MenB, Bexsero, GSK; Trumenba, Pfizer). These vaccines are composed of proteins also found on the surface of the bacteria. Both MenB vaccines are approved by the Food and Drug Administration for use in persons 10 through 25 years of age.

MPSV4 and MenACWY provide no protection against serogroup B disease and meningococcal serogroup B vaccines (MenB) provide no protection against serogroup A, C, W, or Y disease. For protection against all 5 serogroups of meningococcus, it is necessary to receive MenACWY or MPSV4 and MenB.

### **Which individuals in risk groups are recommended to be vaccinated against meningococcal serogroup B disease?**

CDC’s Advisory Committee on Immunization Practices (ACIP) recommends routine MenB vaccination of the following individuals in certain risk groups:

- People age 10 years and older who have functional or anatomic asplenia
- People age 10 years and older who have persistent complement component deficiency, including people taking eculizumab (Soliris)
- People age 10 years and older who are at risk during an outbreak caused by a vaccine serogroup, such as on a college campus
- Microbiologists who work with meningococcus bacteria in a laboratory

Administration of MenB vaccine in persons older than 25 years of age is an off-label use. Clinicians may choose to use vaccines off-label if they believe it would be of benefit to their patients.

### **Which individuals are recommended to be vaccinated against meningococcal serogroup B disease who are not in risk groups?**

ACIP recommends that a MenB vaccine series may be administered to people 16 through 23 years of age with a preferred age of vaccination of 16 through 18 years. This Category B recommendation gives clinicians an opportunity to discuss the value of MenB vaccination with their patients to make a decision together about the individual’s need or desire for the vaccine based on risks, benefits, and wish for protection from the disease. Because it is a Category B recommendation, MenB vaccination is covered by the Vaccines for Children Program for anyone who is eligible. Under the Affordable Care Act, private insurance must also cover the costs of both Category A and B recommended vaccines.

### **What is the difference between a Category A and Category B recommendation?**

A Category A recommendation is made for all persons in an age- or risk-factor-based group. The meningococcal conjugate vaccine recommendation for all preteens at 11–12 years of age is an example of a Category A recommendation. A Category B recommendation does not apply to everyone, but in

the context of a clinician-patient interaction, vaccination may be found to be appropriate for a person as noted above for MenB vaccination of healthy adolescents.

### **Does the Affordable Care Act (ACA) require health plans (non-grandfathered) to provide benefit coverage on Category B recommended vaccines?**

Yes. ACA requires coverage of vaccines with both Category A and B recommendations. The Vaccines for Children Program also includes vaccines with a Category A and B recommendations.

### **Should college students be vaccinated against meningococcal B disease?**

Although several small meningococcal serogroup B disease outbreaks have occurred on college campuses since 2013, college students in general are not at higher risk of meningococcal B disease than persons of the same age who are not college students. Consequently, ACIP does not routinely recommend MenB vaccination for college students. However, college students may choose to receive MenB vaccine to reduce their risk of serogroup B meningococcal disease.

### **Should international travelers receive both meningococcal conjugate vaccine and meningococcal serogroup B vaccine?**

Travelers are not considered to be a group at increased risk for serogroup B meningococcal disease and are not recommended to receive serogroup B vaccine. Meningococcal conjugate vaccine (MenACWY) continues to be recommended for certain international travelers (residents of and travelers to sub-Saharan Africa and the Hajj in Saudi Arabia).

### **What is the schedule for administering MenB vaccine?**

Bexsero is a 2-dose series with dose #2 given at least 1 month after dose #1. Trumenba is either a 2-dose series with doses adminis-

CONTINUED ON THE NEXT PAGE ►

tered at least 6 months apart or a 3-dose series with dose #2 and dose #3 administered 2 and 6 months after dose #1. The ACIP recommends that persons at increased risk of meningococcal serogroup B disease (complement component deficiency, functional or anatomic asplenia, at risk during an outbreak of meningococcal B disease and microbiologists) receive either the 2-dose Bexsero series or the 3-dose Trumenba series. Persons not at increased risk (such as healthy adolescents and young adults) can receive either the 2-dose Bexsero series or the 2-dose Trumenba series.

**What is the least amount of time allowable between doses (minimum intervals) when administering either of the MenB vaccines?**

Neither ACIP nor the CDC meningococcal subject matter experts have addressed this issue. So we must assume that the routinely recommended intervals are also the minimum intervals (see previous question). It is important to use these intervals when scheduling doses. However, if these intervals are violated, CDC recommends that the dose can be counted and does not need to be repeated.

**Can the MenB series be completed with a different MenB brand from the one the series was begun with?**

No. You may not switch MenB vaccines in order to complete a series. The series must be started and completed with the same MenB brand.

**I have a patient who was given Trumenba in August. Two months later she was given a dose of Bexsero. How should I proceed with her MenB vaccination series? We stock both vaccines.**

Since the ACIP meningococcal serogroup B vaccine recommendations state that the same vaccine must be used for all doses in the MenB series, the clinician needs to complete a series with one or the other vaccine. If a non-high risk person has already received

1 dose of Bexsero and 1 of Trumenba, then pick a brand and finish a recommended schedule with that brand. Ignore the extra dose of the other product that was already administered. If you choose to use Bexsero, it should be separated from the previous dose of Bexsero by one month. If you choose to use Trumenba, it should be separated from the previous dose of Trumenba by 6 months.

**We have a 1-year-old with congenital asplenia. He already received a series of meningococcal conjugate vaccine. Should we also give him MenB vaccine?**

Use of either meningococcal serogroup B vaccine in persons younger than age 10 years is off-label in the U.S. There is currently no ACIP recommendation for use of this vaccine for this age group. However, Bexsero brand meningococcal B vaccine has been studied in children and is approved for children as young as 2 months of age by the European Medicines Agency (the European version of the U.S. Food and Drug Administration). It is routinely recommended for infants in the United Kingdom (see [www.nhs.uk/conditions/vaccinations/pages/meningitis-b-vaccine.aspx](http://www.nhs.uk/conditions/vaccinations/pages/meningitis-b-vaccine.aspx) for details). A clinician may choose to use a vaccine off-label if, in their opinion, the benefit of the vaccine exceeds the risk from the vaccine. Product information for Bexsero can be found on the European Medicines Agency website at [www.ema.europa.eu/ema](http://www.ema.europa.eu/ema). These doses may not be covered by insurance.

**Can meningococcal conjugate (MenACWY) and MenB vaccines be given at the same visit?**

Yes. Meningococcal conjugate and MenB vaccines can be given at the same visit or at any time before or after the other.

**Which groups of patients should receive a booster dose of MenB vaccine after completion of the series?**

ACIP does not currently recommend booster doses of MenB vaccine for any group.

**By what route should meningococcal B vaccines be administered?**

MenB vaccines are given by the intramuscular route.

**What are the contraindications and precautions to MenB vaccine?**

As with all vaccines, a severe allergic reaction to a vaccine component or a reaction following a prior dose is a contraindication to subsequent doses. The tip caps of the Bexsero pre-filled syringes contain natural rubber latex which may cause allergic reactions in latex-sensitive individuals. The only precaution for administering MenB vaccine is the presence of a moderate or severe acute illness. Vaccination should be deferred until the illness improves.

**What adverse reactions have been reported after MenB vaccine?**

For both MenB vaccines the most common adverse reactions observed in clinical trials were local reactions, including pain at the injection site (83%–85%), erythema and swelling.

**How should MenB vaccines be stored?**

MenB vaccines should be stored refrigerated at 36°F to 46°F (2°C to 8°C). Do not freeze the vaccines. Discard any vaccine that has been exposed to freezing temperature. Protect the vaccine from light.

**REFERENCES**

- CDC. Use of Serogroup B Meningococcal Vaccines in Persons Aged  $\geq 10$  Years at Increased Risk for Serogroup B Meningococcal Disease: Recommendations of the Advisory Committee on Immunization Practices, 2015. *MMWR* 2016;64(No.22):608-12.
- CDC. Use of Serogroup B Meningococcal Vaccines in Adolescents and Young Adults: Recommendations of the Advisory Committee on Immunization Practices, 2015. *MMWR* 2015;64(No.41):1171-6.

# Meningococcal Vaccine Recommendations by Age and Risk Factor for Serogroup B Protection

This document covers MenB vaccine. For information on vaccine that provides protection against meningococcal serogroup A, C, W, and Y disease, see [www.immunize.org/catg.d/p2018.pdf](http://www.immunize.org/catg.d/p2018.pdf).

- Meningococcal serogroup type B vaccines:**
- **Bexsero** (MenB-4C, GlaxoSmithKline)
  - **Trumenba** (MenB-FHbp, Pfizer)

Routine Recommendations for Meningococcal Serogroup B Vaccination	
For teens and young adults ages 16 through 23 years who wish to be vaccinated. The preferred age is 16 through 18 years.	Give either 2 doses of Bexsero 4 weeks apart, or 2 doses of Trumenba on a 0- and 6-month schedule.

Risk-based Recommendations for Persons with Underlying Medical Conditions or Other Risk Factors	
<p>For people ages 10 years or older with</p> <ul style="list-style-type: none"> <li>• persistent complement component deficiencies<sup>1</sup></li> <li>• anatomic or functional asplenia, including sickle cell disease,</li> </ul> <p>For people ages 10 years or older who</p> <ul style="list-style-type: none"> <li>• are present during outbreaks caused by serogroup B<sup>2</sup></li> <li>• have prolonged increased risk for exposure (e.g., microbiologists routinely working with <i>Neisseria meningitidis</i>)</li> </ul>	Give either 2 doses of Bexsero 4 weeks apart, or 3 doses of Trumenba on a 0-, 2-, and 6-month schedule.

**Note: The two brands of meningococcal B vaccine are not interchangeable. The series must be started and completed with the same brand of vaccine.**

**FOOTNOTES**

1. Persistent complement component deficiencies include inherited or chronic deficiencies in C3, C5–C9, properdin, factor D, and factor H, or taking eculizumab (Soliris)
2. Seek advice of local public health authorities to determine if vaccination is recommended.

# Meningococcal B Vaccine Standing Orders Template Now Updated!

Use this 2-page MenB standing orders template to streamline vaccination of adolescents and young adults in your practice setting.

## Standing Orders for Administering Meningococcal B Vaccine to Adolescents and Adults (continued)

- 5 Administer MenB vaccine, 0.5 mL, via the intramuscular (IM) route, according to the table:

TYPE OF VACCINE	AGE GROUP	DOSE	SCHEDULE
Bexsero <sup>1</sup> (MenB-4c, GlaxoSmithKline)	10 years and older	0.5 mL	Two doses, 4 weeks apart <sup>2,3</sup>
Trumenba <sup>1</sup> (MenB-FHbp, Pfizer)	10 years and older	0.5 mL	Two doses at 0 and 6 months <sup>2</sup> Three doses at 0, 1–2, and 6 months <sup>3</sup>

### Notes:

- The two brands of MenB vaccine are not interchangeable. Do not administer the same brand of vaccine.
- The 2-dose schedules of either Bexsero or Trumenba are preferred.
- Either the 2-dose schedule of Bexsero or the 3-dose schedule of Trumenba is preferred for young adults at increased risk for meningococcal disease due to complement deficiencies, anatomical or functional asplenia, or other conditions.

### 6 Document Vaccination

Document each patient's vaccine administration in the medical record. **Medical record:** Record the date the vaccine was administered, the site and route, and the name and title of the person administering the vaccine. If the vaccine was not administered, record the reason (e.g., patient refusal).

**Personal immunization record card:** Record the date of vaccination in the **Immunization Information System (IIS)** or "registry" if available.

### 7 Be Prepared to Manage Medical Emergencies

Be prepared for management of a medical emergency. Review the emergency medical protocol available, as well as the "Vaccine Reactions in Children and Teens," go to [www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/2013-01-10-Vaccine-Reactions-in-Children-and-Teens.pdf](#) and "Vaccine Reactions in Adult Patients," go to [www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/2013-01-10-Vaccine-Reactions-in-Adult-Patients.pdf](#) while they are seated or lying down and vaccinated.

### 8 Report Adverse Events to VAERS

Report all adverse events following the administration of the vaccine to the Vaccine Adverse Event Reporting System (VAERS) at [www.vaers.hhs.gov](#) or call 822-7967.

## Standing Orders Authorization

This policy and procedure shall remain in effect from \_\_\_\_\_ until rescinded or until \_\_\_\_\_, DATE.  
Medical Director's signature \_\_\_\_\_

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009

Standing orders for other vaccines are available at [www.immunize.org/standing-orders](#). NOTE: This standing orders template may be adapted per a practice's discretion without obtaining permission from IAC. As a courtesy, please acknowledge IAC as its source.

## STANDING ORDERS FOR Administering Meningococcal B Vaccine to Adolescents and Adults

### Purpose

To reduce morbidity and mortality from serogroup B meningococcal disease by vaccinating all adolescents and adults who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP).

### Policy

Where allowed by state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess the need for and vaccinate adolescents and adults who meet any of the criteria below.

### Procedure

#### 1 Assess adolescents and adults for need of vaccination against meningococcal serogroup B disease according to the following criteria:

- Age 16 through 23 years who desire to be vaccinated. The ACIP-preferred age is 16 through 18 years.
- Age 10 years and older, including all adults, with
  - Diagnosis of persistent complement component deficiency (e.g., inherited chronic deficiencies in C3, C5–C9, properdin, factor D and factor H) or taking eculizumab (Soliris)
  - Diagnosis of anatomic or functional asplenia (including sickle cell disease)
  - Risk of potential exposure due to an outbreak attributable to serogroup B
  - Microbiologists routinely exposed to isolates of *Neisseria meningitidis*

#### 2 Screen for contraindications and precautions

**Contraindication** – Do not give meningococcal B vaccine to an adolescent or adult who has experienced a serious systemic or anaphylactic reaction to a prior dose of meningococcal B vaccine or to any of its components. For information on vaccine components, refer to the manufacturers' package insert ([www.immunize.org/packageinserts](#)) or go to [www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf](#).

**Precaution** – Moderate or severe acute illness with or without fever

#### 3 Provide Vaccine Information Statements

Provide all patients (or, in the case of minors, their parent, or legal representative) with a copy of the most current federal Vaccine Information Statement (VIS). Provide non-English speaking patients with a copy of the VIS in their native language, if one is available and desired; these can be found at [www.immunize.org/vis](#). (For information about how to document that the VIS was given, see section 6 titled "Document Vaccination.")

#### 4 Prepare to Administer Vaccine

Choose the needle gauge, needle length, and injection site according to the following chart:

GENDER AND WEIGHT OF PATIENT	NEEDLE GAUGE	NEEDLE LENGTH	INJECTION SITE
Female or male less than 130 lbs	22–25	5/8"–1"	Deltoid muscle of arm
Female or male 130–152 lbs	22–25	1"	Deltoid muscle of arm
Female 153–200 lbs	22–25	1–1½"	Deltoid muscle of arm
Male 153–260 lbs	22–25	1–1½"	Deltoid muscle of arm
Female 200+ lbs	22–25	1½"	Deltoid muscle of arm
Male 260+ lbs	22–25	1½"	Deltoid muscle of arm

\* A 5/8" needle may be used in patients weighing less than 130 lbs (<60 kg) for IM injection in the deltoid muscle only if the skin is stretched tight, the subcutaneous tissue is not bunched, and the injection is made at a 90° angle to the skin.

CONTINUED ON THE NEXT PAGE ►

Technical content reviewed by the Centers for Disease Control and Prevention

IMMUNIZATION ACTION COALITION Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](#) • [www.vaccineinformation.org](#)  
[www.immunize.org/catg.d/p3095.pdf](#) • Item #P3095 (11/16)

Visit [www.immunize.org/catg.d/p3095.pdf](#)



# Meningococcal Vaccine Recommendations by Age and Risk Factor for Serogroups A, C, W, or Y Protection

A separate vaccine is needed for protection against meningococcal serogroup B disease.

**MenACWY** = Menactra (Sanofi Pasteur) and Menveo (GlaxoSmithKline)  
**MenACWY-D** = Menactra      **Hib-MenCY** = MenHibrix (GlaxoSmithKline)  
**MenACWY-CRM** = Menveo      **MPSV** = Menomune (Sanofi Pasteur)

Routine Recommendations for Quadrivalent Meningococcal Conjugate Vaccine (MenACWY)	
For preteens age 11 through 12 years	Give dose #1 of 2-dose MenACWY series. (Dose #2 is recommended at age 16 years.)
For teens age 13 through 15 years	Give catch-up dose #1 of 2-dose MenACWY series. (Dose #2 will be due at age 16 years. <sup>1</sup> )
For teens at age 16 years	Give dose #2 of MenACWY. <sup>1</sup> (Separate from dose #1 by at least 8 weeks.)
Catch-up for teens age 17 through 18 years	If dose #2 not given at age 16 years, give dose #2 of MenACWY as catch-up.
Catch-up for teens age 16 through 18 years	If no history of prior vaccination with MenACWY, give 1 dose of MenACWY.
For first year college students, age 19 through 21 years, living in residence halls	If no history of prior vaccination with MenACWY, give 1 dose of MenACWY. If history of 1 dose of MenACWY given when younger than age 16 years, give dose #2 of MenACWY.

Risk-based Recommendations for Persons with Underlying Medical Conditions or Other Risk Factors		
TARGETED GROUP BY AGE/OR RISK FACTOR	PRIMARY DOSE(S)	BOOSTER DOSE(S)
<b>Travelers to or residents of countries where meningococcal disease is hyperendemic or epidemic,<sup>2</sup> people present during outbreaks caused by a vaccine serogroup,<sup>3</sup> and other people with prolonged increased risk for exposure (e.g., microbiologists routinely working with <i>Neisseria meningitidis</i>)</b>		
For age 2 through 6 months	Give 3 doses of MenACWY-CRM or Hib-MenCY, <sup>4</sup> 8 weeks apart, and a 4th dose at 12–15 months. If possible, vaccination should begin at age 2 months.	If risk continues, give initial booster after 3 years followed by boosters every 5 years.
For age 7 through 23 months who have not initiated a series of MenACWY-CRM	Give 2 doses of MenACWY-CRM <sup>5</sup> or HibMenCY <sup>4,6</sup> or, if 9–23 months, MenACWY-D. <sup>7</sup> Separate the 2 doses by at least 12 weeks. <sup>8</sup>	
For age 2 through 55 years	Give 1 dose of MenACWY.	Boost every 5 years with MenACWY. <sup>9,10</sup>
For age 56 years and older	If no previous MenACWY dose and either short-term travel or outbreak-related, give 1 dose of MPSV; all others, give 1 dose of MenACWY.	Boost every 5 years with MenACWY. <sup>10</sup>
<b>People with persistent complement component deficiencies<sup>11</sup></b>		
For age 2 through 6 months	Give 3 doses of MenACWY-CRM or Hib-MenCY, 8 weeks apart, and a 4th dose at 12–15 months. If possible, vaccination should begin at age 2 months.	Give MenACWY booster after 3 years followed by boosters every 5 years thereafter.
For age 7 through 23 months who have not initiated a series of MenACWY-CRM	Give 2 doses of MenACWY-CRM <sup>5</sup> or Hib-MenCY <sup>6</sup> or, if age 9–23 months, MenACWY-D. <sup>7</sup> Separate the 2 doses by at least 12 weeks.	
For ages 2 through 55 years	Give 2 doses of MenACWY, 8 weeks apart.	Boost every 5 years with MenACWY. <sup>9,12</sup>
For age 56 years and older	Give 2 doses of MenACWY, 8 weeks apart.	Boost every 5 years with MenACWY. <sup>12</sup>
<b>People with HIV infection or functional or anatomic asplenia (including sickle cell disease)</b>		
For age 2 through 6 months	Give 3 doses of MenACWY-CRM or Hib-MenCY, 8 weeks apart, and a 4th dose at 12–15 months. If possible vaccination should begin at age 2 months.	Give MenACWY booster after 3 years followed by boosters every 5 years thereafter. <sup>9</sup>
For age 7 through 23 months who have not initiated a series of MenACWY-CRM	Give 2 doses of MenACWY-CRM <sup>5</sup> or Hib-MenCY. <sup>6</sup> Separate the 2 doses by at least 12 weeks. Or, if using MenACWY-D, give dose #1 at least 4 weeks following completion of pneumococcal conjugate vaccine series, and dose #2 at least 12 weeks after dose #1. <sup>7</sup>	
For ages 2 through 55 years	Give 2 doses of MenACWY, 8 weeks apart.	Boost every 5 years with MenACWY. <sup>9,12</sup>
For age 56 years and older	Give 2 doses of MenACWY, 8 weeks apart.	Boost every 5 years with MenACWY. <sup>12</sup>

## FOOTNOTES

- The minimum interval between doses of MenACWY is 8 weeks.
- Prior receipt of Hib-MenCY is not sufficient for children traveling to the Hajj or African meningitis belt as it doesn't provide protection against serogroups A or W.
- Seek advice of local public health authorities to determine if vaccination is recommended.
- Children ages 2 through 18 months who are present during outbreaks caused by serogroups C or Y may be given an age-appropriate series of Hib-MenCY.
- If initiating vaccination with MenACWY-CRM in a child age 7 through 23 months, dose 2 should be given no younger than age 12 months.
- Hib-MenCY is not licensed for use in children age 18 months or older.
- If MenACWY-D is to be administered to a child with increased risk for meningococcal disease, it should be given either before or concomitantly with DTaP.
- If child age 7 through 23 months will enter an endemic area in less than 3 months, give doses as close as 2 months apart.
- If most recent dose given when younger than age 7 years, give booster after 3 years; if given at or after age 7 years, give booster after 5 years; then boost every 5 years thereafter.
- Booster doses are recommended if the person remains at increased risk.
- Persistent complement component deficiencies include C3, C5–C9, properdin, factor D, factor H, or taking Soliris (eculizumab).
- If the person has a history of only 1 dose, give dose 2 at least 8 weeks after dose 1, then boost every 5 years.

Technical content reviewed by the Centers for Disease Control and Prevention

# Standing Orders Templates Updated for Administering MenACWY Vaccines to Adults and Children/Teens

Standing orders for other vaccines are available at [www.immunize.org/standing-orders](http://www.immunize.org/standing-orders).  
NOTE: This standing orders template may be adapted per a practice's discretion without obtaining permission from IAC. As a courtesy, please acknowledge IAC as its source.

## STANDING ORDERS FOR Administering Meningococcal ACWY Vaccine to Adults

### Purpose

To reduce morbidity and mortality from meningococcal disease caused by serotypes A, C, W, or Y by vaccinating all adults who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices.

### Policy

Where allowed by state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess the need for and vaccinate adults who meet any of the criteria below.

### Procedure

#### 1 Assess adults for need of vaccination against meningococcal disease according to the following criteria:

##### Routine meningococcal ACWY vaccination

- First-year college students age 19 through 21 years living in a residence hall who were never vaccinated or who were last vaccinated when younger than age 16 years

##### Risk-based meningococcal ACWY vaccination

- Diagnosis of persistent complement component deficiency (may also be caused by the drug Soliris [eculizumab])
- Diagnosis of anatomic or functional asplenia (including sickle cell disease)
- Diagnosis of human immunodeficiency virus (HIV) infection
- Part of an outbreak attributable to a vaccine serogroup
- Anticipated travel to a country where meningococcal disease is endemic (the "meningitis belt" of sub-Saharan Africa), particularly if contact with the local population will be prolonged
- Employment as a microbiologist with routine exposure to meningococci

#### 2 Screen for contraindications and precautions

**Contraindications** – Do not give MenACWY vaccine to an adult who has had an anaphylactic reaction to a prior dose of the vaccine or to any of its components, refer to the manufacturer's package insert ([www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf](http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf))

**Precaution** – Moderate or severe acute illness with or without fever

#### 3 Provide Vaccine Information Statements

Provide all patients with a copy of the most current federal Vaccine Information Statement (VIS) available at [www.immunize.org/vis](http://www.immunize.org/vis). You must document in the patient's medical record the date of the VIS and the date it was given to the patient. Provide a copy of the VIS in their native language, if one is available and desired; these can be found at [www.immunize.org/vis](http://www.immunize.org/vis).

#### 4 Review the vaccination schedule and criteria for MenACWY and other meningococcal vaccines

For schedule of vaccination of adults with risk factors as identified in the "Meningococcal Vaccination Recommendations by Age and Risk Factor for Adults" found at [www.immunize.org/catg.d/p2018.pdf](http://www.immunize.org/catg.d/p2018.pdf).

IMMUNIZATION ACTION COALITION Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](http://www.immunize.org)

## FOR ADULTS:

[www.immunize.org/catg.d/p3081.pdf](http://www.immunize.org/catg.d/p3081.pdf)

## FOR CHILDREN/TEENS:

[www.immunize.org/catg.d/p3081a.pdf](http://www.immunize.org/catg.d/p3081a.pdf)

Download and use these standing orders templates "as they are," or modify them to suit your work setting.

Standing orders for other vaccines are available at [www.immunize.org/standing-orders](http://www.immunize.org/standing-orders).  
NOTE: This standing orders template may be adapted per a practice's discretion without obtaining permission from IAC. As a courtesy, please acknowledge IAC as its source.

## STANDING ORDERS FOR Administering Meningococcal ACWY Vaccine to Children and Teens

### Purpose

To reduce morbidity and mortality from meningococcal disease caused by serotypes A, C, W, or Y by vaccinating all children and teens who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices.

### Policy

Where allowed by state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess the need for and vaccinate children and teens who meet any of the criteria below.

### Procedure

#### 1 Assess children and teens for need of vaccination against meningococcal disease according to the following criteria:

##### Routine meningococcal ACWY vaccination

- Age 11–12 years who have not received MenACWY at age 10 years or older
- As catch-up for ages 13–15 years who have not received MenACWY at age 10 years or older
- Age 16 years and in need of dose #2
- Ages 17 through 18 years and in need of dose #2 as catch-up
- As catch-up for all unvaccinated teens ages 16 through 18 years
- First-year college students age 19 through 21 years living in a residence hall who were never vaccinated or who were last vaccinated when younger than age 16 years

##### Risk-based meningococcal ACWY vaccination

- Age 2 months and older with diagnosis of persistent complement component deficiency (an immune system disorder, which may also be caused by the drug Soliris [eculizumab]), diagnosis of anatomic or functional asplenia (including sickle-cell disease), or diagnosis of infection with human immunodeficiency virus; children who are part of an outbreak attributable to a vaccine serogroup; or anticipated travel to a country where meningococcal disease is hyperendemic or epidemic (e.g., the "meningitis belt" of sub-Saharan Africa), particularly if contact with the local population will be prolonged

#### 2 Screen for contraindications and precautions

**Contraindications** – Do not give MenACWY vaccine to a child or teen who has a history of a serious allergic reaction (e.g., anaphylaxis) after a previous dose of meningococcal vaccine or to a meningococcal vaccine component. For information on vaccine components, refer to the manufacturer's package insert ([www.immunize.org/packageinserts](http://www.immunize.org/packageinserts)) or go to [www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf](http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf).

**Precaution** – Moderate or severe acute illness with or without fever

#### 3 Provide Vaccine Information Statements

Provide all patients (or, in the case of a minor, their parent or legal representative) with a copy of the most current federal Vaccine Information Statement (VIS) available at [www.immunize.org/vis](http://www.immunize.org/vis). You must document in the patient's medical record or office log, the publication date of the VIS and the date it was given to the patient (parent/legal representative). Provide non-English speaking patients with a copy of the VIS in their native language, if one is available and desired; these can be found at [www.immunize.org/vis](http://www.immunize.org/vis).

#### 4 Prepare to Administer Vaccine

Choose the needle gauge, needle length, and injection site according to the following chart:

AGE OF PATIENT	NEEDLE GAUGE	NEEDLE LENGTH	INJECTION SITE
Adolescents (age 11–21 years)	22–25	5/8"–1"	Deltoid muscle of arm
Children (age 3–10 years)	22–25	5/8"–1"	Deltoid muscle of arm
Toddlers (age 1–2 years)	22–25	1–1½"	Anterolateral thigh muscle
Infants (age 2–12 months)	22–25	1"	Anterolateral thigh muscle

\* A 5/8" needle may be used in patients weighing less than 130 lbs (<60 kg) for IM injection in the deltoid muscle only if the skin is stretched tight, the subcutaneous tissue is not bunched, and the injection is made at a 90-degree angle to the skin.

(PAGE 1 OF 2)

IMMUNIZATION ACTION COALITION Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](http://www.immunize.org) • [www.vaccineinformation.org](http://www.vaccineinformation.org)  
[www.immunize.org/catg.d/p3081a.pdf](http://www.immunize.org/catg.d/p3081a.pdf) • Item #P3081a (12/16)

# Great Resources on [www.Give2MCV4.org](http://www.Give2MCV4.org) to Help Protect Preteens and Teens from Meningococcal A, C, W, Y Disease



- ▶ Meningococcal conjugate vaccine (MCV4) provides safe and effective protection against meningococcal disease caused by serogroups A, C, W, and Y.
- ▶ MCV4 is recommended at ages 11–12 followed by a second (booster) vaccination at age 16.
- ▶ According to CDC’s 2015 National Immunization Survey–Teen, **only 33% of teens had received their recommended booster dose by 17 years of age.**

**Valuable Resource!** Downloadable slide deck and speaker notes for healthcare professionals

[www.Give2MCV4.org](http://www.Give2MCV4.org)

## More Resources

Visit [www.Give2MCV4.org](http://www.Give2MCV4.org) to view the full collection of resources designed to help healthcare professionals improve rates for MCV4 and all recommended adolescent vaccines, including:

**Recommending MCV4: What to Say and How to Say It**  
[www.give2mcv4.org/wp-content/uploads/2015/07/Toolkit-Recommending-MCV4.pdf](http://www.give2mcv4.org/wp-content/uploads/2015/07/Toolkit-Recommending-MCV4.pdf)


**Top 10 Ways to Improve Adolescent Immunization Rates**  
[www.give2mcv4.org/wp-content/uploads/2015/07/Toolkit-Top-10-Ways.pdf](http://www.give2mcv4.org/wp-content/uploads/2015/07/Toolkit-Top-10-Ways.pdf)

**Screening Checklist for Contraindications to HPV, MCV4, MenB, and Tdap**  
[www.immunize.org/catg.d/p4062.pdf](http://www.immunize.org/catg.d/p4062.pdf)

and much more!

**“Dear Colleague” Letter: Call-to-Action from IAC, CDC, and professional societies emphasizing the importance of the second dose of MCV4**  
[www.immunize.org/mcv4letter](http://www.immunize.org/mcv4letter)

**MCV4 YOU'RE NOT DONE IF YOU GIVE JUST ONE**  
**GIVE 2 DOSES** to Strengthen Protection



# Newly Updated Meningococcal Q&As: Download and Copy for Your Patients

## Meningococcal: Questions and Answers

### INFORMATION ABOUT THE DISEASE AND VACCINES

#### What causes meningococcal disease?

Meningococcal disease is caused by the bacterium *Neisseria meningitidis*. This bacterium has at least 13 different subtypes (serogroups). Five of these serogroups, A, B, C, Y, and W, cause almost all invasive disease. The relative importance of these five serogroups depends on geographic location and other factors. In the United States almost all meningococcal disease is caused by serogroups B, C, and Y. Each serogroup accounts for about one third of reported cases.

#### How does meningococcal disease spread?

The disease is spread person-to-person through the exchange of respiratory and throat secretions (e.g., by coughing, kissing, or sharing eating utensils). Meningococcal bacteria can't live for more than a few minutes outside the body, so the disease is not spread as easily as the common cold or influenza.

#### How long does it take to show signs of meningococcal disease after being exposed?

The incubation period of meningococcal disease is 3 to 4 days, with a range of 2 to 10 days. Meningococcal bacteria can make a person extremely ill by infecting the blood (septicemia) or by infecting the fluid of the spinal cord and around the brain (meningitis). Because this disease progresses quickly, it is important to be diagnosed and start treatment as soon as possible.

#### What are the symptoms of meningococcal disease?

The most common symptoms are high fever, chills, lethargy, and a rash. If meningitis is present, the symptoms will also include headache and neck stiffness (which may not be present in infants); seizures may also occur. In overwhelming meningococcal infections, shock, coma, and death can follow within several hours, even with appropriate medical treatment.

#### How serious is meningococcal disease?

Meningococcal disease caused by any serogroup is very serious. About 10 to 15% of people with meningococcal disease die even with appropriate antibiotic treatment. Of those who recover, up to 20% suffer

from some serious after-effects, such as permanent hearing loss, limb loss, or brain damage.

#### How is meningococcal disease diagnosed?

The diagnosis is made by taking samples of blood and spinal fluid from a person who is sick. The spinal fluid is obtained by performing a lumbar puncture, where a needle is inserted into the lower back. Any bacteria found in the blood or spinal fluid is grown in a medical laboratory and identified.

Meningococcal disease is uncommon in the United States, and the symptoms can be mistaken for other illnesses, which unfortunately can lead to delayed diagnosis and treatment.

#### Can't meningitis be caused by a virus too?

Yes. The word "meningitis" refers to inflammation of the tissues covering the brain and spinal cord. This inflammation can be caused by viruses and fungi, as well as bacteria. Viral meningitis is the most common type. It has no specific treatment but is usually not as serious as meningitis caused by bacteria.

#### Is there a treatment for meningococcal disease?

Meningococcal disease can be treated with antibiotics. It is critical to start treatment early.

#### How common is meningococcal disease in the United States?

Fewer than 700 cases of meningococcal disease were reported each year since 2010 in the United States. An estimated average 80 deaths from meningococcal disease occurred each year in the United States since 2010.

The disease is most common in children younger than 5 years (particularly children younger than age 1 year), people age 16–21 years, and people age 65 years and older.

#### What people are at special risk for meningococcal disease?

For all meningococcal serogroups risk factors include age, having a damaged or missing spleen, persistent

CONTINUED ON THE NEXT PAGE ►

Technical content reviewed by the Centers for Disease Control and Prevention  
Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org  
www.immunize.org/catg.d/p4210.pdf • Item #P4210 (11/16)

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org  
www.immunize.org/catg.d/p4210.pdf • Item #P4210 (11/16)

#### Who should not receive meningococcal vaccine?

These groups should not receive either type of meningococcal vaccine:

- People who have had a serious allergic reaction to a previous dose of either meningococcal vaccine or to one of the vaccine components. The packaging of some meningococcal vaccines may contain latex. Information on the contents of each vaccine is included with each vaccine.
- People who are moderately or severely ill.

#### Can a pregnant woman get meningococcal vaccine?

Studies of vaccination with MPSV4 during pregnancy have not documented adverse effects among either pregnant women or newborns. Post-licensure safety data suggest no concerns with the safety of MenACWY during pregnancy. Pregnancy is not considered to be a contraindication to either MPSV4 or MenACWY. Although experience with MenB vaccines is limited they have not been shown to be detrimental to a pregnant woman or fetus.

#### Can the vaccine cause meningococcal disease?

No. Only the *Neisseria meningitidis* bacterium can cause meningococcal disease. Meningococcal vaccines contain only the sugar capsule or capsule protein of the microbe.

and microbiologists receive either the 2-dose Bexsero series or the 3-dose Trumenba series. Persons not at increased risk (such as healthy adolescents and young adults) can receive either the 2-dose Bexsero series or the 2-dose Trumenba series. Booster doses of MenB vaccine following the initial series are currently not recommended, including for people with no spleen or persistent complement component deficiency.

#### How soon after their first MenACWY dose should people who remain at risk for meningococcal disease be vaccinated again?

The time between the primary (initial) dose(s) of MenACWY and the first booster varies. Children who received their primary MenACWY dose(s) before their seventh birthday should get their first booster 3 years after their primary dose(s). Children who received their primary MenACWY dose(s) at or after age 7 years and all adults should get MenACWY boosters 5 years after their primary dose(s).

#### What are the side effects of this vaccine?

Up to about half of people who get meningococcal vaccines have mild side effects, such as redness or pain where the shot was given. These symptoms usually last for one or two days and are more common after MenACWY than after MPSV4. A small percentage of people who receive the vaccine develop a fever. Severe reactions, such as a serious allergic reaction, are very rare. More than 60,000 persons have received MenB vaccines during clinical trials or for outbreak control on college campuses. The most common side effect was pain at the injection site, which was reported by about 80% of recipients. The Vaccine Adverse Event Reporting System (VAERS) and other vaccine safety systems will carefully monitor MenB vaccine safety as they do for other U.S.-licensed vaccines.

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org  
www.immunize.org/catg.d/p4210.pdf • Item #P4210 (11/16)



continued) page 2 of 4

(An immune system people taking the cupation as a micro- with meningococcal

for meningococcal serogroup B. These s where meningococ- tain countries in ffection, and college /see question below for having a previous household, having being exposed to second-hand).

#### occur in other parts

throughout the world, f Africa known as the responsible for most jib-Saharan Africa. he United States.

#### an once?

ifferent serogroups y other bacteria

children and adults (e.g., who are in the same school or church) aren't usually considered exposed unless they have had very close contact with the infected person (e.g., kissing or sharing a glass).

In addition to the antibiotic treatment, vaccination may be recommended for people 2 months of age and older if the person's infection is caused by meningococcus serogroup A, C, Y, or W-135, which are contained in 3 of the 5 meningococcal vaccines available in the United States.

#### What meningococcal vaccines are available in the United States?

There are 3 types of meningococcal vaccine available in the United States. Vaccines for meningococcal serogroups A, C, W and Y represent 2 of the 3 types. One is composed of polysaccharide (sugar molecules) from the surface of the meningococcal bacteria. The other meningococcal vaccines (MenACWY) are those in which the polysaccharide is chemically bonded ("conjugated") to a protein. These produce better protection and are more effective in young children than the original polysaccharide vaccine. A third type are vaccines for meningococcal serogroup B (MenB), which are composed of proteins also found in the surface of the bacteria. No type of vaccine contains live meningococcal bacteria.

Meningococcal polysaccharide or conjugate vaccines provide no protection against serogroup B disease and

Meningococcal Vaccines Licensed in U.S.				
TRADE NAME	TYPE OF VACCINE	SEROGROUPS INCLUDED	YEAR LICENSED	APPROVED AGES
Menomune	Polysaccharide	A, C, W, Y	1981	2 years and older
Menactra	Conjugate	A, C, W, Y	2005	9 months–55 years*
Menveo	Conjugate	A, C, W, Y	2010	2 months–55 years**
Trumenba	Protein	B	2014	10–25 years†
Bexsero	Protein	B	2015	10–25 years†

\*may be given to people age 56 years or older  
\*\*may be given to people age 26 years or older

to a person with ed by being started tely (ideally within nosed). This is usu- contacts and children sery school. Older

MenB vaccines provide no protection against serogroup A, C, W or Y disease. For protection against all 5 serogroups of meningococcus it is necessary to receive meningococcal conjugate or polysaccharide (that contain serogroup A, C, W, and Y) and a MenB vaccine.

CONTINUED ON THE NEXT PAGE ►

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org  
www.immunize.org/catg.d/p4210.pdf • Item #P4210 (11/16)

This 4-page Q&A for patients is ready for you to hand out in your medical setting.

Visit [www.immunize.org/catg.d/p4210.pdf](http://www.immunize.org/catg.d/p4210.pdf)

ed) page 3 of 4

PSV4) is for the upper MenACWY or the deldoid tenB vaccines

ccine? CWY and to receive only for cer-

ups: 18 years re a damaged

re persistent immun- ned by the during an

hey are or will residential hall de in or travel ca as well as ccal vaccine Saudi Arabia,

istered to persons 16 through 23 years of age with a preferred age of vaccination of 16 through 18 years. This permissive (Category B) recommendation allows the clinician to make a MenB vaccine recommendation based on the risk and benefit for the individual patient.

#### Should college students be vaccinated against meningococcal disease?

The MenACWY vaccine is recommended for previously unvaccinated first-year college students, age younger than 22 years, who are or will be living in a residence hall. Some colleges and universities require incoming freshmen and others to be vaccinated with MenACWY; some may also require that a dose of MenACWY have been given since the age of 16 years. MenACWY may be available from the college health service.

Although several small meningococcal B disease outbreaks have occurred on college campuses since 2013, college students in general are not at higher risk of meningococcal B disease than persons of the same age who are not college students. Consequently, ACIP does not routinely recommend MenB vaccination for college students. However, college students may choose to receive MenB vaccine to reduce their risk should a meningococcal B disease outbreak occur.

#### How many doses of meningococcal vaccine are needed?

For MenACWY vaccines the number of doses recommended depends on the age when the vaccine is given and the presence of certain medical conditions or risk factors. All adolescents should be vaccinated with one dose of MenACWY at ages 11 or 12 years and with a booster dose at age 16 years. All teens who were vaccinated with MenACWY at ages 13 through 15 years need a booster dose at age 16 through 18 years (at least 8 weeks after the first dose). First-year college students younger than 22 years who are or will be living in a residential hall should get a MenACWY booster dose if their previous dose was given before age 16 years. People ages 2 months and older who have certain risk factors such as no spleen or a damaged spleen, or persistent complement component deficiency (an immune system disorder which may also be caused by the drug Soliris [eculizumab]), may need more than one dose. In addition, vaccinated people who remain at risk, such as people without a spleen, people with HIV infection, microbiologists who work with meningococcus, or

CONTINUED ON THE NEXT PAGE ►

► For IAC's ready-to-print Q&As about all vaccine-preventable diseases, visit [www.immunize.org/handouts/vaccine-questions.asp](http://www.immunize.org/handouts/vaccine-questions.asp)

# Here's More Meningococcal Vaccine Information for Your Patients

An easy-to-read fact sheet for teens and adults

## Protect yourself from meningococcal disease... Get vaccinated!

### What is meningococcal disease?

Meningococcal disease can be a life-threatening illness. It is caused by bacteria that can infect the blood, brain, and spinal cord. People often call it meningitis.

### How do you catch it?

Meningococcus bacteria are spread through upper respiratory droplets, like saliva (kissing, living in close quarters). You can catch meningitis from a person who looks healthy.



### Is it serious?

Yes! Meningococcal disease may respond to antibiotics, but quick medical attention is extremely important. Even with proper treatment, 10–15% of people with meningococcal disease die. Of the people who survive, as many as 20% suffer from some serious complication, such as loss of an arm or leg, brain damage, or permanent hearing loss.

Ask your healthcare provider if you need this vaccine!

Meningitis can cause shock, coma, and death within hours of the first symptom.

### Am I at risk?

The disease most often strikes older teens and young adults. If you travel internationally or have certain medical conditions, you may also be at risk.

### How can I protect myself from meningitis?



Vaccination is the best way to prevent meningococcal disease.

There are different types of meningococcal vaccines: MenACWY conjugate and MenB (serogroup B). MenACWY is given to preteens and teens beginning at age 11–12 years. A second dose is needed at age 16. A MenB series can also be given to teens and young adults starting at age 16.

People with certain medical conditions should get vaccinated, and some should receive booster doses throughout life. Travelers to certain countries should also get vaccinated.

Download from IAC's website:

[www.immunize.org/catg.d/p4410.pdf](http://www.immunize.org/catg.d/p4410.pdf)

► For more information, visit [www.vaccineinformation.org](http://www.vaccineinformation.org)

For other vaccine handouts in this series, visit [www.immunize.org/vaccine-summaries](http://www.immunize.org/vaccine-summaries)



Technical content reviewed by the Centers for Disease Control and Prevention  
Saint Paul, Minnesota • [www.immunize.org](http://www.immunize.org)  
[www.immunize.org/catg.d/p4410.pdf](http://www.immunize.org/catg.d/p4410.pdf) • Item #P4410 (9/16)

► For IAC's easy-to-read fact sheets about all vaccine-preventable diseases (including Spanish translations), visit [www.immunize.org/handouts/vaccine-summaries.asp](http://www.immunize.org/handouts/vaccine-summaries.asp)

NEWLY UPDATED!

# Which Vaccines Do I Need Today?

This checklist helps you determine which vaccines your adult patients need.

Download and copy this screening questionnaire for your patients to fill out.

YOUR NAME \_\_\_\_\_ DATE OF BIRTH \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_ TODAY'S DATE \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_  
month / day / year month / day / year

## Which Vaccines Do I Need Today?

Vaccines are an important part of helping you stay healthy. Which of these recommended vaccines do you need? Check the boxes that apply to you, and then talk this over with your healthcare provider.

### Influenza ("flu") vaccine

- I have not had my flu vaccine yet this season (*early fall through late spring*).

### Pneumococcal ("pneumonia") vaccines [Pneuvax 13 [PCV13] and Pneumovax 23 [PPSV23]]

I am **age 65 or older** and:

- I have never received any pneumonia vaccine (or I don't remember if I have).
- I have received only 1 pneumonia vaccine since I turned 65.
- I received 1 or 2 doses of pneumonia vaccine before I turned 65, and it's now been since I received my last dose.

I am **younger than age 65** and:

- I have never received any pneumonia vaccine AND at least one of the following:
  - I smoke cigarettes and I am age 19 years or older.
  - I have a chronic disease of the heart, lung (including asthma, if I am age 19 years or older), kidneys, or I have sickle cell disease.
  - I have diabetes or alcoholism.
  - I have a weakened immune system due to cancer, Hodgkin's disease, leukemia, myeloma, kidney failure, HIV/AIDS or receiving radiation therapy or taking an immune system.
  - I live in a nursing home or other long-term care facility.
- I have had an organ or bone marrow transplant.
- I have had my spleen removed or have had a cochlear (inner ear) implant or have a provider that I have leaking spinal fluid.

### Tetanus, diphtheria, and pertussis ("whooping cough")-containing vaccine (e.g., DTaP)

- I have never received Tdap vaccine (or I don't remember if I have).
- I have not received at least 3 tetanus- and diphtheria-containing shots.
- I have received at least 3 tetanus- and diphtheria-containing shots in my lifetime more than 10 years since I received the last one.
- I am pregnant (and I am in my late second or third trimester of my pregnancy) and I did not receive a Tdap vaccine during this pregnancy.

### Measles, mumps, rubella (MMR) vaccine

- I am a woman thinking about a future pregnancy and don't know if I'm immune.
- I am a healthcare worker. I have received 1 MMR (or I don't remember if I have received more than 1). I do not have a lab-confirmed report showing that I am immune to measles, mumps, and/or rubella. I was born in 1957 or later and:
  - I have never received MMR vaccine (or I don't remember if I have).
  - I have received only 1 MMR and
    - I am entering college or another type of school after high school.
    - I am planning on traveling outside the U.S.<sup>1</sup>

Which Vaccines Do I Need Today? (continued) page 2 of 3

#### Human papillomavirus (HPV) vaccination

I have not completed a series of HPV shots and

- I am a woman age 26 or younger.
- I am a man
  - age 21 or younger.
  - age 22 through 26 and at least one of the following applies to me:
    - I want to be protected from HPV.
    - I have a weakened immune system due to infection (including HIV), disease, or medications.
    - I have sex with men.
- I am now older than age 26 and have not completed the HPV vaccine series I began when I was age 26 or younger.

#### Hepatitis A vaccine

- I want to be vaccinated to avoid getting hepatitis A and spreading it to others.
- I might have been exposed to hepatitis A virus within the past 2 weeks.
- I received 1 dose of hepatitis A vaccine in the past, but I have not received the second dose (or I don't remember if I have).
- I have not received hepatitis A vaccine in the past (or I don't remember if I have) and at least one of the following applies to me:
  - I travel (or plan to travel) in countries where hepatitis A is common.<sup>1,2</sup>
  - I have (or will have) contact with a child within 60 days of the child's adoption from a country where hepatitis A is common.<sup>2</sup>
  - I am a man who has sex with men.
  - I use street drugs.
  - I have chronic liver disease.
  - I have a blood clotting factor disorder.
  - I work with hepatitis A virus in a research laboratory or with primates infected with hepatitis A virus.

#### Hepatitis B vaccine

- I want to be vaccinated to avoid getting hepatitis B and spreading it to others.
- I am age 18 or younger and I have not begun or completed the series of hepatitis B shots (or I don't remember if I have).
- I have received at least one dose of hepatitis B in the past, but I have not completed the series of hepatitis B shots (or I don't remember if I have).
- I have not received or completed the series of hepatitis B shots (or I don't remember if I have) and at least

Which Vaccines Do I Need Today? (continued) page 3 of 3

#### Varicella ("chickenpox") vaccine

- I was born before 1980 and I am a healthcare worker or foreign-born and I don't remember if I've ever had chickenpox disease.
- I was born in 1980 or later and I have never had chickenpox disease or received the vaccine (or I don't remember if I have).
- I have received one dose of varicella vaccine, but I'm not sure if I have received more than one dose.

#### Meningococcal ("meningitis") type A, C, W, Y vaccine (MenACWY [MCV4])

- I am age 18 or younger and have never received any meningococcal vaccines (or I don't remember if I have).
- I am age 21 or younger and
  - I have not had a meningococcal shot (MenACWY) since before my 16th birthday and I am (or will be) a college student living in a residence hall.
  - I have not had a meningococcal shot (MenACWY) in the past 5 years and I am entering college.
- I have sickle cell disease.
- My spleen isn't working or has been removed.
- I have a persistent complement component deficiency.
- I have HIV infection.
- I have a risk of exposure due to an outbreak caused by serogroup A, C, W, or Y.
- I am a microbiologist who is routinely exposed to isolates of *Neisseria meningitidis*.
- I was vaccinated more than 5 years ago and I continue to be at risk due to travel, illness, or occupation.

#### Meningococcal ("meningitis") type B vaccine (MenB)

- I am age 16-23 with no specific risk factor and would like to be protected from this disease.
- I have a risk of exposure due to an outbreak caused by serogroup B.
- I have sickle cell disease.
- My spleen isn't working or has been removed.
- I have a persistent complement component deficiency.

#### Zoster ("shingles") vaccine

- I am age 60 or older and have never received a shingles vaccine (or I don't know if I have).

#### Haemophilus influenzae type b ("Hib") vaccine

- My spleen has been removed, or I am scheduled to have it removed ("splenectomy").
- I have received a stem cell transplant.

#### Travel vaccines

- I am planning on traveling outside the U.S.<sup>1,3</sup> (Discuss this with your provider.)

**FOOTNOTES**

1. Call your local travel clinic to find out if additional vaccines are recommended.
2. Countries where hepatitis A is common include all countries other than the U.S., Western Europe, Canada, Japan, Australia, and New Zealand.
3. Areas with high rates of hepatitis B include Africa, China, Korea, Southeast Asia including Indonesia and the Philippines, South and Western Pacific Islands, interior Amazon Basin, certain parts of the Caribbean (i.e., Haiti and the Dominican Republic), and the Middle East except Israel. Areas with moderate rates include South Central and Southeast Asia, Israel, Japan, Eastern and Southern Europe, Russia, and most of Central and South America.
4. Most adults from moderate- or high-risk areas of the world do not know their hepatitis B status. All patients from these areas need hepatitis B blood tests to determine if they have been previously infected. The first hepatitis B shot can be given during the same visit as the blood tests but only after the blood is drawn.

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org  
www.immunize.org/catg.d/p4036.pdf • Item #P4036 (1/16)

▶ Visit [www.immunize.org/catg.d/p4036.pdf](http://www.immunize.org/catg.d/p4036.pdf)



# Use This Checklist to Maximize Protection of Your Valuable Vaccine Supply

## Checklist for Safe Vaccine Storage and Handling

Are you doing everything you should to safeguard your vaccine supply? Review this list to see where you might make improvements in your vaccine management practices. Check each listed item with either **YES** or **NO**.

### Establish Storage and Handling Policies

- YES**  **NO** 1. We have designated a primary vaccine coordinator and at least one alternate coordinator to be in charge of vaccine storage and handling at our facility.
- YES**  **NO** 2. Both the primary and alternate vaccine coordinator(s) have completely reviewed either CDC Vaccine Storage & Handling Toolkit ([www.cdc.gov/vaccines/hcp/admin/storage/toolkit/storage-handling.pdf](http://www.cdc.gov/vaccines/hcp/admin/storage/toolkit/storage-handling.pdf)) or equivalent training materials offered by our state or local health department's immunization program.
- YES**  **NO** 3. We have detailed, up-to-date, written standard operating procedures for general vaccine management including procedures for routine activities and an emergency vaccine retrieval and storage outages and other problems. Our procedures are based on CDC's Vaccine Storage & Handling Toolkit and/or on instruction from our state or local health department's immunization program.
- YES**  **NO** 4. We review these policies with all staff annually and with new staff, including temporary staff who are hired.

### Log In New Vaccine Shipments

- YES**  **NO** 5. We maintain a vaccine inventory log that we use to document the following:
- YES**  **NO** a. Vaccine name and number of doses received
  - YES**  **NO** b. Date we received the vaccine
  - YES**  **NO** c. Condition of vaccine when we received it
  - YES**  **NO** d. Vaccine manufacturer and lot number
  - YES**  **NO** e. Vaccine expiration date

### Use Proper Storage Equipment

- YES**  **NO** 6. We store vaccines in separate, sealed, hold-style combination unit, we store vaccines in a separate stand-alone unit.
- YES**  **NO** 7. We store vaccines in units with emergency power backup.
- YES**  **NO** 8. We never store any vaccines in a refrigerator or freezer compartment inside a stand-alone unit.
- YES**  **NO** 9. We use only calibrated thermometers and are calibrated every 1 month or as the manufacturer's suggested time interval.
- YES**  **NO** 10. We have planned back-up storage for our vaccines.

\*Certificate of Calibration Testing ("Report of Call to a laboratory with accreditation from the International Laboratory Accreditation Cooperation (ILAC) Mutual Recognition Arrangement (MRA).")



Saint Paul, Minnesota • 651-647-9009

Are you doing everything you should to safeguard your vaccine supply? Review this 3-page checklist to see where you might make improvements.

## Checklist for Safe Vaccine Storage and Handling (continued)

page 2 of 3

### Ensure Optimal Operation of Storage Units

- YES**  **NO** 11. We have a "Do Not Unplug" sign (e.g., [www.immunize.org/catg.d/p2090.pdf](http://www.immunize.org/catg.d/p2090.pdf)) next to the electrical outlets for the refrigerator and freezer and a "Do Not Stop Power" warning label (e.g., [www.immunize.org/catg.d/p2091.pdf](http://www.immunize.org/catg.d/p2091.pdf)) by the circuit breaker for the electrical outlets. Both signs include emergency contact information.
- YES**  **NO** 12. We perform regular maintenance on our vaccine storage units to assure optimal functioning. For example, we keep the units clean, dusting the coils and cleaning beneath the units as recommended by the manufacturer.

### Maintain Correct Temperatures

- YES**  **NO** 13. We always keep at least one accurate calibrated thermometer (+/-0.5°C [+/-1°F]) with the vaccines in the refrigerator and a separate calibrated thermometer with the vaccines in the freezer.
- YES**  **NO** 14. We use a thermometer that
- YES**  **NO** a. uses an active display to provide continuous monitoring information.
  - YES**  **NO** b. is digital and has a detachable probe that has been buffered against sudden temperature changes by being immersed in a vial filled with liquid (e.g., glycol, ethanol, glycerin), loose media (e.g., sand, glass beads), or a solid block of material (e.g., aluminum, Teflon®).
  - YES**  **NO** c. includes an alarm for out-of-range temperatures.
  - YES**  **NO** d. has a digital data logger that indicates current, minimum, and maximum temperatures.
  - YES**  **NO** e. can measure temperatures within +/-0.5°C (+/-1°F).
  - YES**  **NO** f. has a low-battery indicator.
- YES**  **NO** 15. We maintain the refrigerator temperature at 2–8°C (36–46°F), and we aim for 5°C (40°F).
- YES**  **NO** 16. We maintain the freezer temperature between -50°C and -15°C (-58°F and +5°F).
- YES**  **NO** 17. We set the thermostat for the refrigerator and the freezer at the factory-set or midpoint temperatures.
- YES**  **NO** 18. We keep extra containers of water in the refrigerator (e.g., in the door and/or on the floor of the unit where the vegetable bins were located) to help maintain cool temperatures. We keep ice packs, ice-filled containers, or frozen water bottles in the freezer to help maintain cold temperatures and to have frozen water bottles available for conditioning in the event of an emergency.
- ### Maintain Daily Temperature Logs
- YES**  **NO** 19. On days when our practice is open, we visually inspect the vaccine storage unit twice a day (first thing in the morning and right before our facility closes) and document refrigerator and freezer temperatures on the appropriate log. (See selections at [www.immunize.org/clinic/storage-handling.asp](http://www.immunize.org/clinic/storage-handling.asp).)
- YES**  **NO** 20. We document the minimum and maximum temperature readings in the refrigerator and freezer once each day, preferably in the morning.
- YES**  **NO** 21. We consistently record temperatures on the log either in Fahrenheit or Celsius. We never mix temperature scales when we record our temperatures.
- YES**  **NO** 22. If the temperature log prompts us to insert an "x" by the temperature that's preprinted on the form, we do not attempt to write in the actual temperature.
- YES**  **NO** 23. We follow the directions on the temperature log to call appropriate personnel if the temperature in a storage unit goes out of range.
- YES**  **NO** 24. If out-of-range temperatures occur in the unit, we complete the Vaccine Storage Troubleshooting Record ([www.immunize.org/catg.d/p3041.pdf](http://www.immunize.org/catg.d/p3041.pdf)) to document actions taken when the problem was discovered and what was done to prevent a recurrence of the problem.

CONTINUED ON THE NEXT PAGE ▶

Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](http://www.immunize.org) • [www.vaccineinformation.org](http://www.vaccineinformation.org)  
[www.immunize.org/catg.d/p3035.pdf](http://www.immunize.org/catg.d/p3035.pdf) • Item #P3035 (11/16)

## Checklist for Safe Vaccine Storage and Handling (continued)

- YES**  **NO** 25. Trained staff (other than staff designated to handle vaccines) are responsible for monitoring the temperature logs on file for at least 1 year.
- YES**  **NO** 26. We keep the temperature logs on file for at least 1 year.

### Store Vaccines Correctly

- YES**  **NO** 27. We post signs (e.g., [www.immunize.org/catg.d/p2092.pdf](http://www.immunize.org/catg.d/p2092.pdf)) that indicate which vaccines should be stored in the refrigerator and which should be stored in the freezer.
- YES**  **NO** 28. We do not store any food or drink in any vaccine storage unit.
- YES**  **NO** 29. We store vaccines in the middle of the refrigerator (not near the top shelf). We do not store vaccines in the door.
- YES**  **NO** 30. We have removed all vegetable and deli bin empty areas.
- YES**  **NO** 31. If we must use a combination refrigerator-freezer, we do not place vaccines in the freezer (often near the top shelf). In general, we do not place water bottles in this location.
- YES**  **NO** 32. We check vaccine expiration dates and rotate the earliest expiration dates are located close to the front of the unit.
- YES**  **NO** 33. We store vaccines in their original packaging.

### Take Emergency Action As Needed

- YES**  **NO** 34. In the event that vaccines are exposed to improper storage conditions:
- YES**  **NO** a. We restore proper storage conditions as soon as possible and move it to a unit where it can be stored before discussing the circumstances with the appropriate vaccine manufacturers.
  - YES**  **NO** b. We follow the Vaccine Storage Troubleshooting Record instructions for taking appropriate action, such as the length of time the vaccine was exposed to room temperature, as well as taking an inventory of the unit.
  - YES**  **NO** c. We contact our clinic supervisor or other appropriate personnel, our state/local health department and/or the appropriate vaccine manufacturers for consultation about whether the exposed vaccine can still be used.
  - YES**  **NO** d. We address the storage unit's mechanical or electrical problems according to guidance from the unit's manufacturer or a qualified repair service.
  - YES**  **NO** e. In responding to improper storage conditions, we do not make frequent or large changes in thermostat settings. After changing the setting, we give the unit at least a day to stabilize its temperature.
  - YES**  **NO** f. We do not use exposed vaccines until our state/local health department's immunization program or the vaccine manufacturer has confirmed that the vaccine is acceptable for use. We review this information with our clinic medical director before returning the vaccine to our supply. If the vaccine is not acceptable for use, we follow our state/local health department instructions for vaccine disposition.

If we answer **YES** to all of the above, we give ourselves a pat on the back! If not, we assign someone to implement needed changes!


Immunization Action Coalition • Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](http://www.immunize.org) • [www.vaccineinformation.org](http://www.vaccineinformation.org)  
[www.immunize.org/catg.d/p3035.pdf](http://www.immunize.org/catg.d/p3035.pdf) • Item #P3035 (11/16)

▶ Visit [www.immunize.org/catg.d/p3035.pdf](http://www.immunize.org/catg.d/p3035.pdf)



# Vaccine Handling Tips

REMEMBER: Improperly stored or outdated vaccines won't protect your patients!



## Manage vaccine inventories.

Inventory your vaccine supplies at least monthly and before placing an order. Expired vaccine must never be used and it becomes “cash in the trash!”

## Always use the vaccine with the soonest expiration date first.

Move vaccine with the soonest expiration date to the front of the storage unit and mark it to be used first. These actions help ensure it will be picked up first by someone selecting vaccine from the unit.

## Store vaccine appropriately.†

Place vaccines in refrigerator or freezer immediately upon receiving shipment. Keep vaccine vials in their original packaging. Place vaccine in clearly labeled wire baskets or other open containers with a 2–3" separation between baskets and 4" from wall of unit. Separate or clearly mark vaccines to distinguish those that were supplied from your state's Vaccines for Children program (or other state-funded source) from those that were privately purchased. Do not store vaccines in the door or on the floor of the unit.

## Stabilize temperatures.

Store ice packs in the freezer and large jugs of water in the refrigerator along with the vaccines. This will help maintain a stable, cold temperature in case of a power failure or if the refrigerator or freezer doors are opened frequently or are accidentally left open. Because frequent opening of either the refrigerator or freezer door can lead to temperature variations that could affect vaccine efficacy, you should not store food or beverages in the refrigerator or freezer.

## Safeguard the electrical supply to the refrigerator.

Make sure the refrigerator and freezer are plugged into outlets in a protected area where they cannot be disconnected accidentally. Label the refrigerator, freezer, electrical outlets, fuses, and circuit breakers on the power circuit with information that clearly identifies the perishable nature of vaccines and the immediate steps to be taken in case of interruption of power.‡ If your building has auxiliary power, use the outlet supplied by that system.

\*MMR may be stored in either the freezer or the refrigerator.

† Refer to package insert for specific instructions on the storage of each vaccine. If you have questions about the condition of the vaccine upon arrival, immediately place the vaccine in recommended storage, mark it “do not use,” and then call your state health department or the vaccine manufacturer(s) to determine whether the potency of the vaccine(s) has been affected. For other questions, call the immunization program at your state or local health department.


‡ For easy help with labeling units and power supplies, see IAC signs “Do Not Unplug Refrigerator or Freezer” ([www.immunize.org/catg.d/p2090.pdf](http://www.immunize.org/catg.d/p2090.pdf)) and “Do Not Stop Power to Circuit Breaker” ([www.immunize.org/catg.d/p2091.pdf](http://www.immunize.org/catg.d/p2091.pdf)). For guidance on steps to take during a power interruption, see IAC's “Emergency Response Worksheet” ([www.immunize.org/catg.d/p3051.pdf](http://www.immunize.org/catg.d/p3051.pdf)).

# These products are available for purchase from the Immunization Action Coalition

Laminated adult and child/teen immunization schedules — Order one of each for every exam room

▶ To order, visit [www.immunize.org/shop](http://www.immunize.org/shop), or use the order form on page 20.

Coming in March 2017: The ACIP/AAFP/ACOG/ACNM-approved schedule for adults (8-sided) and the ACIP/AAP/AAFP-approved immunization schedule for people ages 0 through 18 years (8-sided). Both are laminated and washable for heavy-duty use, complete with essential footnotes, and printed in color for easy reading.



Schedules: \$7.50 each  
Quantity discounts are available.



Wallet-sized immunization record cards for all ages: For adults, for children and teens, and for a lifetime!

Record Cards: \$45/box  
Now you can give any patient a permanent vaccination record card designed specifically for their age group: adult, child and teen, or lifetime. These brightly colored cards are printed on durable rip-, smudge-, and water-proof paper. Each box contains 250 cards.

▶ To order, visit [www.immunize.org/shop](http://www.immunize.org/shop), or use the order form on page 20.

Quantity discounts are available. To receive sample cards, contact us: [admininfo@immunize.org](mailto:admininfo@immunize.org)



Training Video: “Immunization Techniques — Best Practices with Infants, Children, and Adults”

The California Department of Public Health, Immunization Branch, updated its award-winning training video, “Immunization Techniques: Best Practices with Infants, Children, and Adults.” The 25-minute DVD can be used to train new employees and to refresh the skills of experienced staff on administering injectable, oral, and nasal-spray vaccines to children, teens, and adults.

DVD: \$17 each  
Quantity discounts are available.


▶ To order, visit [www.immunize.org/shop](http://www.immunize.org/shop), or use the order form on page 20.

For healthcare settings in California, contact your local health department immunization program for a free copy.

## The Vaccine Handbook: A Practical Guide for Clinicians (“The Purple Book”) by Gary S. Marshall, MD

During my more than 25 years in the field of immunization education, I have not seen another book that is so brimming with state-of-the-science information. — DEBORAH L. WEXLER, MD, Executive Director, IAC

Purchase *The Vaccine Handbook* (560 pages) from IAC at [www.immunize.org/vaccine-handbook](http://www.immunize.org/vaccine-handbook).  
\$29.95 + shipping • Discount pricing available.



# Help Us Increase Immunization Rates!



The Immunization Action Coalition (IAC) is the nation's premier source of immunization information for healthcare professionals. Your contribution directly funds our websites, publications, email news service, and handouts that improve the delivery of vaccines and motivate patients to get immunized!



## I want to support IAC!

### Here is my contribution

I am a  new  renewing contributor

\$100  \$50  \$75  \$250

\$500  \$750  \$1,000

other \$ \_\_\_\_\_

IAC is a 501(c)(3) charitable organization and your contribution is tax deductible to the fullest extent of the law.

► **DONATE BY FAX** (651) 647-9131

► **DONATE BY MAIL**

Immunization Action Coalition  
2550 University Avenue West, Suite 415 N  
Saint Paul, MN 55114

► **DONATE ONLINE**

[www.immunize.org/support](http://www.immunize.org/support)

NAME/TITLE \_\_\_\_\_

ORGANIZATION \_\_\_\_\_

ADDRESS \_\_\_\_\_  
Check one: This is my  organization address  home address

CITY/STATE/ZIP \_\_\_\_\_

PHONE \_\_\_\_\_ EMAIL \_\_\_\_\_

**Method of payment**  Check made out to Immunization Action Coalition

Purchase order

Visa  Mastercard  Am Express  Discover

Credit Card #

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

CV Code #\*

--	--	--	--	--

\* CV Code # – the additional 3- or 4-digit number on your credit card

Expiration date (month/year) 

--	--	--	--

--	--	--	--

## Honor a colleague, friend, or relative with your contribution!

A gift to IAC is a thoughtful way to honor or remember someone. When you make an honorary or memorial gift, we will send your honoree an acknowledgement that expresses our sincere thanks for the support your gift provides.

IAC may identify me to the honoree as the source of this contribution.

IAC may acknowledge the amount of the gift to the honoree.

A separate letter will be sent to you (the donor) for your tax records.

My contribution is  in honor of  in memory of:

NAME OF HONOREE \_\_\_\_\_

OCCASION (birthday, anniversary, memorial, etc.) \_\_\_\_\_

Please send a letter of acknowledgement to:

NAME (if different) \_\_\_\_\_

INSTITUTION (if applicable) \_\_\_\_\_

ADDRESS \_\_\_\_\_

CITY/STATE/ZIP \_\_\_\_\_

# Order Essential Immunization Resources from IAC

Immunization record cards for all: for adults, for children and teens, for a lifetime!

Immunization record cards give healthcare professionals a way to help patients maintain a permanent record of their vaccinations. Having one's own vaccination record is handy for patients when they enter daycare, kindergarten, or college; change healthcare providers; or travel abroad. ■ The Immunization Action Coalition offers three record cards: adult, child and teen, and lifetime. Each is designed for

a specific age group and lists all vaccines recommended for people in that age group. Sized to fit in a wallet, each is brightly colored to stand out and is printed on durable rip-, smudge-, and water-proof paper. ■ To order record cards or any of our other essential immunization resources, print out and mail or fax the form below, or place your order online at [www.immunize.org/shop](http://www.immunize.org/shop).

It's convenient to shop IAC online at [www.immunize.org/shop](http://www.immunize.org/shop)

## ■ The Vaccine Handbook: A Practical Guide for Clinicians ("The Purple Book") by Gary Marshall, MD

Fifth edition • 2015 • 560 pages • \$29.95 + shipping

Order online at [www.immunize.org/vaccine-handbook](http://www.immunize.org/vaccine-handbook)



## Order Essential Immunization Resources

### ■ Coming March 2017: Laminated 2016 U.S. Immunization Schedules (details p. 20; call for discounts on bulk orders)

Qty.	1-4 copies—\$7.50 each; 5-19 copies—\$5.50 each	Amt.
_____	R2008 Child/teen immunization schedules .....	\$ _____
_____	R2009 Adult immunization schedules .....	\$ _____

### ■ DVD – Immunization Techniques: Best Practices with Infants, Children, and Adults (details p. 20; call for discounts on bulk orders)

1-9 copies—\$17 each; 10-24 copies—\$10.25 each; 25-49 copies—\$7 each

_____	D2021 Immunization Techniques: Best Practices with Children/Teens/Adults	\$ _____
-------	--	----------

### ■ Patient Immunization Record Cards – for children and teens, for adults, and for a lifetime! (all are wallet-sized; details p. 20; call for discounts on bulk orders)

250 cards/box; 1 box—\$45; 2 boxes—\$40 each; 3 boxes—\$37.50 each; 4-7 boxes—\$34.50 each

_____	R2003 Child/teen immunization record cards .....	\$ _____
_____	R2005 Adult immunization record cards .....	\$ _____
_____	R2004 Lifetime immunization record cards .....	\$ _____

**Total for Purchases \$** \_\_\_\_\_

### ■ Make a Charitable Contribution

I am a  new  renewing contributor.

\$100  \$250  \$50  \$35 other: \$ \_\_\_\_\_

IAC is a 501(c)(3) charitable organization and your contribution is tax deductible to the fullest extent of the law.

**Total for Purchases and Contribution \$** \_\_\_\_\_

## How to Place an Order

**By Credit Card:** Order easily online at our secure shopping cart at [www.immunize.org/shop](http://www.immunize.org/shop).

**By Check, Purchase Order, or Credit Card:** Print out this page, fill out the necessary information, and

**Fax this page to:** (651) 647-9131 or

**Mail this page to:** Immunization Action Coalition  
2550 University Avenue West, Suite 415 North  
Saint Paul, MN 55114

**Our federal ID# is 41-1768237.**

**For Questions or International Orders:** Contact us by phone at (651) 647-9009 or email [admininfo@immunize.org](mailto:admininfo@immunize.org)

**Thank you** for your support of the Immunization Action Coalition. We depend on you!

**Method of payment:**  Check enclosed (payable to Immunization Action Coalition)

Purchase order # \_\_\_\_\_

Visa  Mastercard  Am. Express  Discover

Card #

Expiration Date mo/yr   CV Code #\*

\*The CV Code is the Credit Verification Code, the additional 3- or 4-digit number on your credit card.

Name/Title \_\_\_\_\_

Organization \_\_\_\_\_

Shipping address (Check one: This is my  organization address  home address)

City/State/Zip \_\_\_\_\_

Telephone \_\_\_\_\_

Email address \_\_\_\_\_

It's convenient to shop IAC online at [www.immunize.org/shop](http://www.immunize.org/shop)