


# Laminated U.S. Immunization Schedules

Purchase IAC's laminated versions of the 2017 U.S. immunization schedules for adults as well as for children/teens (0–18 years old). Both are laminated and washable for heavy-duty use, complete with essential footnotes, and printed in color for easy reading.

Schedules are \$7.50 each; quantity discounts are available. More information and discount pricing options are available online at [www.immunize.org/shop/laminated-schedules.asp](http://www.immunize.org/shop/laminated-schedules.asp) or see the order form on page 18.



## Recommended Immunization Schedules for Children and Adolescents Ages 0 through 18 Years, United States, 2017

Also included: Guide to Contraindications and Precautions to Commonly Used Vaccines

## Recommended Adult Immunization Schedule, United States, 2017

Also included: Guide to Contraindications and Precautions to Commonly Used Vaccines in Adults

To order additional copies of this laminated document, visit the Immunization Action Coalition's website for healthcare professionals at [www.immunize.org/hcp](http://www.immunize.org/hcp), call (851) 647-9009, or email [advis@immunize.org](mailto:advis@immunize.org).



St. Paul, Minnesota • (851) 647-9009 • [www.immunize.org](http://www.immunize.org) • fax

To order additional copies of this laminated document, visit the Immunization Action Coalition's website for healthcare professionals at [www.immunize.org/hcp](http://www.immunize.org/hcp), call (851) 647-9009, or email [advis@immunize.org](mailto:advis@immunize.org).



St. Paul, Minnesota • (851) 647-9009 • [www.immunize.org](http://www.immunize.org) • [www.vaccineinformation.org](http://www.vaccineinformation.org) • [advis@immunize.org](mailto:advis@immunize.org) • fax (612) 928-5171

Recommended Adult Immunization Schedule—United States, 2017		Footnotes		Table 1. Guide to Contraindications and Precautions to Commonly Used Vaccines in Adults <sup>1, 2, 3</sup>	
<p><b>1. Minimum vaccinees</b></p> <p>Adults aged 18 years and older who have not received a complete series of adult immunizations should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>	<p><b>2. Recommended vaccinees</b></p> <p>Adults aged 18 years and older who have received a complete series of adult immunizations should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>	<p><b>3. Recommended vaccinees</b></p> <p>Adults aged 18 years and older who have received a complete series of adult immunizations and are at high risk for infection should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>	<p><b>4. Recommended vaccinees</b></p> <p>Adults aged 18 years and older who have received a complete series of adult immunizations and are at high risk for infection should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>	<p><b>5. Recommended vaccinees</b></p> <p>Adults aged 18 years and older who have received a complete series of adult immunizations and are at high risk for infection should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>	<p><b>6. Recommended vaccinees</b></p> <p>Adults aged 18 years and older who have received a complete series of adult immunizations and are at high risk for infection should receive the following vaccines:</p> <ul style="list-style-type: none"> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> <li><b>MMWR</b> (Measles, Mumps, Rubella)</li> <li><b>Tdap</b> (Tetanus, Diphtheria, Pertussis)</li> <li><b>Polio</b> (Poliovirus)</li> <li><b>Shingles</b> (Herpes zoster)</li> <li><b>Hepatitis A</b> (Hepatitis A virus)</li> <li><b>Hepatitis B</b> (Hepatitis B virus)</li> <li><b>Pneumococcal</b> (Pneumococcus)</li> </ul>