

Meningococcal: Questions and Answers

INFORMATION ABOUT THE DISEASE AND VACCINES

What causes meningococcal disease?

Meningococcal disease is caused by the bacterium *Neisseria meningitidis*. This bacterium has at least 13 different subtypes (serogroups). Five of these serogroups, A, B, C, Y, and W, cause almost all invasive disease. The relative importance of these five serogroups depends on geographic location and other factors. In the United States almost all meningococcal disease is caused by serogroups B, C and Y. Each serogroup accounts for about one third of reported cases.

How does meningococcal disease spread?

The disease is spread person-to-person through the exchange of respiratory and throat secretions (e.g., by coughing, kissing, or sharing eating utensils). Meningococcal bacteria can't live for more than a few minutes outside the body, so the disease is not spread as easily as the common cold or influenza.

How long does it take to show signs of meningococcal disease after being exposed?

The incubation period of meningococcal disease is 3 to 4 days, with a range of 2 to 10 days. Meningococcal bacteria can make a person extremely ill by infecting the blood (septicemia) or by infecting the fluid of the spinal cord and around the brain (meningitis). Because this disease progresses quickly, it is important to be diagnosed and start treatment as soon as possible.

What are the symptoms of meningococcal disease?

The most common symptoms are high fever, chills, lethargy, and a rash. If meningitis is present, the symptoms will also include headache and neck stiffness (which may not be present in infants); seizures may also occur. In overwhelming meningococcal infections, shock, coma, and death can follow within several hours, even with appropriate medical treatment.

How serious is meningococcal disease?

Meningococcal disease caused by any serogroup is very serious. About 15% of people with meningococcal disease die even with appropriate antibiotic treatment. Of those who recover, up to 20% suffer from some

serious after-effects, such as permanent hearing loss, limb loss, or brain damage.

How is meningococcal disease diagnosed?

The diagnosis is made by taking samples of blood and spinal fluid from a person who is sick. The spinal fluid is obtained by performing a lumbar puncture, where a needle is inserted into the lower back. Any bacteria found in the blood or spinal fluid is grown in a medical laboratory and identified.

Meningococcal disease is uncommon in the United States, and the symptoms can be mistaken for other illnesses, which unfortunately can lead to delayed diagnosis and treatment.

Can't meningitis be caused by a virus too?

Yes. The word "meningitis" refers to inflammation of the tissues covering the brain and spinal cord. This inflammation can be caused by viruses and fungi, as well as bacteria. Viral meningitis is the most common type; it has no specific treatment but is usually not as serious as meningitis caused by bacteria.

Is there a treatment for meningococcal disease?

Meningococcal disease can be treated with antibiotics. It is critical to start treatment early.

How common is meningococcal disease in the United States?

Fewer than 500 cases of meningococcal disease were reported each year since 2010 in the United States. In 2017, a total of 310 cases were reported and 40 died.

The disease is most common in children younger than 5 years (particularly children younger than age 1 year), people age 16–21 years, and people age 65 years and older.

What people are at special risk for meningococcal disease?

For all meningococcal serogroups risk factors include age, having a damaged or missing spleen, persistent complement component deficiency (an immune system

CONTINUED ON THE NEXT PAGE ►

disorder which may also occur in people taking the drug Soliris [eculizumab]), and occupation as a microbiologist in a laboratory that works with meningococcal isolates.

Certain groups are at increased risk for meningococcal serogroups A, C, Y, and W but not serogroup B. These risk factors include travel to places where meningococcal disease is common (such as certain countries in Africa and in Saudi Arabia), HIV infection, and college freshmen who live in a dormitory (see question below for more on college students). Other risk factors for serogroups A, C, Y and W include having a previous viral infection, living in a crowded household, having an underlying chronic illness, and being exposed to cigarette smoke (either directly or second-hand).

Does meningococcal disease occur in other parts of the world?

Meningococcal disease occurs throughout the world, but is more common in the area of Africa known as the “meningitis belt.” Serogroup A is responsible for most of the meningococcal disease in sub-Saharan Africa. This serogroup is uncommon in the United States.

Can you get meningitis more than once?

Yes. Meningitis can be caused by different serogroups of the meningococcal bacterium, by other bacteria such as *Streptococcus* and *Haemophilus*, as well as by viruses and fungi. Being vaccinated against *Neisseria meningitidis* or having had the disease will not protect you against meningitis from other bacteria or viruses.

If a child is diagnosed with meningococcal disease, can anything be done to protect the other children with whom he has contact?

Individuals who have been exposed to a person with bacterial meningitis can be protected by being started on a course of antibiotics immediately (ideally within 24 hours of the patient being diagnosed). This is usually recommended for household contacts and children attending the same day care or nursery school. Older children and adults (e.g., who are in the same school or church) aren’t usually considered exposed unless they have had very close contact with the infected person (e.g., kissing or sharing a glass).

In addition to the antibiotic treatment, vaccination may be recommended for people 2 months of age and older if the person’s infection is caused by meningococcus serogroup A, C, Y, or W, which are contained in 4 of the 5 meningococcal vaccines available in the United States.

What meningococcal vaccines are available in the United States?

There are 2 types of meningococcal vaccine available in the United States. One type of vaccine (MenACWY) contains the surface polysaccharides of meningococcal serogroups A, C, W and Y chemically bonded (conjugated) to a protein. This vaccine is recommended for all adolescents at 11–12 years and a second dose at 16 years. A second type are vaccines for meningococcal serogroup B (MenB), which are composed of proteins also found in the surface of the bacteria. No type of vaccine contains live meningococcal bacteria.

MenACWY vaccines provide no protection against serogroup B disease and MenB vaccines provide no protection against serogroup A, C, W or Y disease. For protection against all 5 serogroups of meningococcus it is necessary to receive a MenACWY and a MenB vaccine.

| Meningococcal Vaccines Licensed in U.S. | | | | |
|---|-----------------|---------------------|---------------|--------------------|
| TRADE NAME | TYPE OF VACCINE | SEROGROUPS INCLUDED | YEAR LICENSED | APPROVED AGES |
| Menactra | Conjugate | A, C, W, Y | 2005 | 9 months–55 years* |
| Menveo | Conjugate | A, C, W, Y | 2010 | 2 months–55 years* |
| Trumenba | Protein | B | 2014 | 10–25 years† |
| Bexsero | Protein | B | 2015 | 10–25 years† |

*may be given to people age 56 years or older
 †may be given to people age 26 years or older

How is this vaccine given?

MenACWY vaccines are given in a leg muscle of a young child or the deltoid (arm) muscle of an older child or adult. MenB vaccines are given in the deltoid muscle.

Who should get the meningococcal vaccine?

Certain groups should receive both MenACWY and MenB vaccines. Others are recommended to receive MenACWY only.

CONTINUED ON THE NEXT PAGE ►

MenACWY is recommended for these groups:

- All children and teens, ages 11 through 18 years
- People age 2 months and older who have a damaged or missing spleen
- People age 2 months and older who have persistent complement component deficiency (an immune system disorder which may also be caused by the drug Soliris [eculizumab]), or are at risk during an outbreak caused by a vaccine serogroup
- People with HIV infection.
- People who are or will be a first-year college student living in a residential facility
- People age 2 months and older who reside in or travel to certain countries in sub-Saharan Africa as well as to other countries for which meningococcal vaccine is recommended (e.g., travel to Mecca, Saudi Arabia, for the annual Hajj).
- People working with meningococcus bacteria in laboratories

MenB is recommended for these groups:

- People age 10 years and older who have a damaged or missing spleen
- People age 10 years and older who have persistent complement component deficiency (an immune system disorder which may also be caused by the drug Soliris [eculizumab]), or are at risk during an outbreak caused by a vaccine serogroup
- People working with meningococcus bacteria in laboratories

MenB vaccines are not routinely recommended for all adolescents or college students. However, the Centers for Disease Control and Prevention (CDC) recommends that a MenB vaccine series may be administered to persons 16 through 23 years of age with a preferred age of vaccination of 16 through 18 years. This permissive recommendation allows the clinician to make a MenB vaccine recommendation based on the risk and benefit for the individual patient.

Should college students be vaccinated against meningococcal disease?

The MenACWY vaccine is recommended for previously unvaccinated first-year college students who are or will be living in a residence facility. Some colleges and universities require incoming freshmen and others to be vaccinated with MenACWY; some may also require that

a dose of MenACWY have been given since the age of 16 years. MenACWY may be available from the college health service.

Although several small meningococcal B disease outbreaks have occurred on college campuses since 2013, college students in general are not at higher risk of meningococcal B disease than persons of the same age who are not college students. Consequently, CDC does not routinely recommend MenB vaccination for college students. However, college students may choose to receive MenB vaccine to reduce their risk should a meningococcal B disease outbreak occur.

How many doses of meningococcal vaccine are needed?

For MenACWY vaccines the number of doses recommended depends on the age when the vaccine is given and the presence of certain medical conditions or risk factors. All adolescents should be vaccinated with one dose of MenACWY at age 11 or 12 years and with a booster dose at age 16 years. All teens who were vaccinated with MenACWY at age 13 through 15 years need a booster dose at age 16 through 18 years (at least 8 weeks after the first dose). First-year college students who are or will be living in a residential facility should get a MenACWY booster dose if their previous dose was given before age 16 years. People ages 2 months and older who have certain risk factors such as no spleen or a damaged spleen, or persistent complement component deficiency (an immune system disorder which may also be caused by the drug Soliris [eculizumab]), may need more than one dose. In addition, vaccinated people who remain at risk, such as people without a spleen, people with HIV infection, microbiologists who work with meningococcus, or those who travel repeatedly to parts of Africa, should receive a booster dose of MenACWY every 5 years.

The CDC recommends that persons at increased risk of meningococcal B disease (complement component deficiency, functional or anatomic asplenia, at risk during an outbreak of meningococcal B disease and microbiologists) receive either the 2-dose Bexsero series or the 3-dose Trumenba series. Persons not at increased risk (such as healthy adolescents and young adults) can receive either the 2-dose Bexsero series or the 2-dose Trumenba series. Booster doses of MenB vaccine following the initial series are currently not recommended, including for people with no spleen or persistent complement component deficiency.

CONTINUED ON THE NEXT PAGE ►

How soon after their first MenACWY dose should people who remain at risk for meningococcal disease be vaccinated again?

The time between the primary (initial) dose(s) of MenACWY and the first booster varies. Children who received their primary MenACWY dose(s) before their seventh birthday should get their first booster 3 years after their primary dose(s). Children who received their primary MenACWY dose(s) at or after age 7 years and all adults should get MenACWY boosters 5 years after their primary dose(s).

What are the side effects of this vaccine?

Up to about half of people who get MenACWY vaccines have mild side effects, such as redness or pain where the shot was given. These symptoms usually last for one or two days. A small percentage of people who receive the vaccine develop a fever. Severe reactions, such as a serious allergic reaction, are very rare.

More than 60,000 persons have received MenB vaccines during clinical trials or for outbreak control on college campuses. The most common side effect was pain at the injection site, which was reported by about 80% of recipients. The Vaccine Adverse Event Reporting System (VAERS) and other vaccine safety systems will carefully monitor MenB vaccine safety as they do for other U.S.-licensed vaccines.

How effective is this vaccine?

Based on antibody studies and comparison with an older meningococcal vaccine, MenACWY is believed to be at least 85% effective.

Because of the low incidence of serogroup B meningococcal disease, MenB vaccine efficacy estimates were based on demonstration of an immune response after

vaccination. From 63% to 88% of recipients of a full series of MenB vaccine develop a protective level of antibody against representative strains of serogroup B meningococcus.

Who should not receive meningococcal vaccine?

These groups should not receive either type of meningococcal vaccine:

- People who have had a serious allergic reaction to a previous dose of either meningococcal vaccine or to one of the vaccine components. The packaging of some meningococcal vaccines may contain latex. Information on the contents of each vaccine is included with each vaccine.
- People who are moderately or severely ill.

Can a pregnant woman get meningococcal vaccine?

Post-licensure safety data suggest no concerns with the safety of MenACWY during pregnancy. Pregnancy is not considered to be a contraindication to MenACWY. Healthcare personnel should administer this vaccine to high-risk pregnant women. Although experience with MenB vaccines is limited, they have not been shown to be detrimental to a pregnant woman or fetus. Pregnancy is a precaution to MenB vaccine. Healthcare personnel can administer this vaccine to high-risk pregnant women.

Can the vaccine cause meningococcal disease?

No. Only the *Neisseria meningitidis* bacterium can cause meningococcal disease. Meningococcal vaccines contain only the sugar capsule or capsule protein of the microbe.