
## Use These Standing Orders Templates for Administering Influenza Vaccination in Your Healthcare Setting

Download these standing orders and use them "as is," or modify them to suit your work setting.

STANDING ORDERS FOR

Administering Influenza Vaccine to Adults



STANDING ORDERS FOR Administering Influenza Vaccine to Children and Adolescents To reduce morbidity and mortality from influenza by vaccinating all children and adolescents who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices (ACIP). Where allowed by state law, standing orders enable eligible nurses and other healthcare professionals (e.g., pharmacists) to assess the need for vaccination and to vaccinate children and adolescents who meet any of the criteria below. NOTE: Live attenuated influenza vaccine (LAIV4; FluMist), is not recommended by CDC's Advisory Committee on Immunization Practices for use in the U.S. during the 2016–17 influenza season. Because LAIV4 is still a licensed vaccine that might be available and that some providers might elect to use, for informational purposes, reference is made to previous recommendations for its use. Procedure 1 Assess Children and Adolescents for Need of Vaccination against influenza Assess Children and Addressens for Inveted of Vaccination against himberida

All children and teens 6 months of age and older are recommended to receive influenza vaccination each year.

A second dose of influenza vaccine is recommended 4 weeks or more after the first dose for children age 6 months through 8 years if they have not received 2 doses in previous years (not necessarily in the same season). 2 Screen for Contraindications and Precautions Screen for Contraindications and Precautions

Contraindications for use of all influenza vaccines

Do not give influenza vaccine to a child or adolescent who has experienced a serious systemic or anaphylactic reaction to a prior dose of the vaccine or to any of its components. For a list of vaccine components, refer to the pinkbook/downloads/appendices/8/excipient-table-2.pdf. Contraindications only for use of five attenuated influenza vaccine (LAIV4; FluMist, nasal spray)

Do not give live attenuated influenza vaccine (LAIV4, nasal spray) to a child or adolescent who:

• is nearmant is pregnant
is younger than age 2 years
is age 2 through 4 years who has experienced wheezing or asthma within the past 12 months, based on a healthcare provider's statement
has immunosuppression (including that caused by medications or HIV)
is age 2 through 17 years and is on long-term aspirin or salicylate-containing therapy
is age 2 through 17 years and is on long-term aspirin or salicylate-containing therapy
48 hours or will possibly receive them within 14 days after vaccination
provides care for a severely immunosuppressed person who requires a protective environment Precautions for use of all influenza vaccines

Moderate or severe acute illness with or without fever

History of Guillain-Barré syndrome within 6 weeks of a previous influenza vaccination Precautions for use of LAIV only

Age 5 years or older with asthma older with asthma

"dical conditions (e.g., other chronic lung diseases, chronic cardiovascular disease (excluding sion), chronic renal or hepatic disease, hematologic disease, neurologic disease, and metacluding diabetes mellitus) ENTS WITH EGGS ALLERGY: People with egg allergy of any severity can receive any licensed

Standing Orders for Administering Influenza Vaccine to Children and Adolescents

ON Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org

www.immunize.org (catg.d/p3074a.pdf • item #93074a [9])

www.immunize.org/catg.d/p3074a.pdf

Standing Orders for Administering Influenza Vaccine to Adults www.immunize.org/catg.d/p3074.pdf

Additional standing orders templates for all routinely recommended vaccines are available at www.immunize.org/standing-orders